

Nonstandard Analysis of Graphs

F. Javier Thayer

February 1, 2008

Abstract

This paper shows certain classes of metric length spaces characterized by volume growth properties of balls can be viewed as graphs with infinitesimal edges. Our approach is based on nonstandard analysis.

1 Introduction

Intuitively, it is tempting to view a cube in $\mathbb{R}^n$ as a discrete lattice with infinitesimal edges. A similar infinitesimal picture suggests itself with other fractal spaces such as the Sierpinski gasket or the Sierpinski sponge (see [21], [8].) This paper shows certain classes of metric spaces characterized by a polynomial growth condition on the volume of balls ([5],[3]) can be viewed as a part of a graph with infinitesimal edges. Our approach uses nonstandard analysis in two ways: first, nonstandard analysis makes it possible to do infinitesimal rescaling of graph distances in a completely straightforward way, and second, we use the well-known connection between nonstandard counting measures and countably additive measures established in [19] and [14] to associate Hausdorff measure on metric spaces to counting measures on graphs. The main result is Theorem 10.5 which exhibits an Ahlfors regular length space up to Lipschitz equivalence as a part of a hyperfinite graph in which the edges have infinitesimal length.

The structure of the paper is as follows: the next two sections consist mainly of reference material, particularly on nonstandard analysis and a summary of results in nonstandard measure theory. In particular, in Section 2 we give a definition of polynomial growth for functions defined on intervals with hyperreal endpoints. Note that for functions which are defined on an interval whose endpoints are real numbers, polynomial growth reduces to boundedness on that interval.

In Section 5, we introduce the main concept of the paper, namely that of polynomial growth for hyperfinite graphs. The basic idea is as follows: Consider the shortest-path metric between nodes. If the cardinality of balls as a function of the radius behaves polynomially in some interval, we say the graph has polynomial growth on that interval. Note the notion of polynomial growth for finite graphs is vacuous. Along with the shortest-distance metric, we consider rescalings of that metric and using nonstandard analysis we have a lot of

leeway on how we choose the rescaling parameter. In particular, if we choose an infinitesimal rescaling, we get a graph in which nodes may be at an infinitesimal distance from each other. Now collapse the graph, identifying nodes that are infinitely close to each other. The resulting object is a metric space, which in the polynomial growth case has a volume growth property called Ahlfors regularity as shown in Theorem 5.4. The remainder of the paper shows that the converse (suitably stated) also holds for the class of length spaces introduced by Gromov in [10].

We note that the relation between Gromov's geometric ideas and nonstandard analysis is not new. Gromov himself uses ultraproducts in a construction he calls the asymptotic cone over a metric space. An explicit connection between nonstandard analysis and asymptotic geometry has been established in the beautiful paper [24]. The authors of that paper used this connection to provide a different proof of Gromov's theorem that a finitely generated group of polynomial growth has a nilpotent subgroup of finite index. However, the relation between polynomial growth of graphs and volume growth seems to be new.

2 Preliminaries

We will consider *extended metric spaces* in which the metric is allowed to assume the value $+\infty$. Note that this generalization does not introduce any new topological behavior. Given an extended metric space (X, d) , define an equivalence relation for pairs x, y of X by $d(x, y) < \infty$. We will call the equivalence classes of this relation the *bounded components* of (X, d) . The bounded components of (X, d) are open and closed sets.

Definition 2.1 *An extended metric space (X, d) is a length space iff for every pair $x, y \in X$ with $d(x, y) < \infty$, there is an isometric map $f : [0, d(x, y)] \rightarrow X$ with $f(0) = x, f(d(x, y)) = y$.*

This is a specialization of Gromov's definition (see [10], Définition 1.7), although by Théorème 1.10 of [10] the two definitions are equivalent for complete locally compact metric spaces. For example, if X is a geodesically complete Riemannian manifold, by the Hopf-Rinow theorem [3], Theorem 1.9, (X, d) is a length space. The following result is Théorème 1.10, (i) of [10]. See also the first chapter of [11]:

Proposition 2.2 *If (X, d) is a complete locally compact length space, all closed balls $\overline{B}(a, r)$ with $r < \infty$ are compact.*

2.1 Ball Borel Structure

If (X, d) is an extended metric space, the smallest σ -algebra containing the open balls $B(a, r)$ with $r < \infty$ is the *ball Borel structure*. If (X, d) is separable, then since every open set in X is a countable union of open balls, the ball Borel

structure is identical to the usual Borel structure generated by the open sets. In general, the ball Borel structure has fewer sets than the Borel structure. For instance, if (X, d) is an uncountable discrete metric space (3.2.5 of [6]), the only balls in X are $\emptyset$, X and singletons. Thus the ball Borel sets of X are precisely those sets which are countable or have countable complements whereas any subset of X is open and so any subset of X is Borel.

If (X, d) is a metric space, and $A \subseteq X$, then the ball Borel structure on (A, d) may not be the same as the ball Borel structure of (X, d) relativized to A . For example, let (A, d) be an uncountable discrete metric space, $X = A \cup \{z\}$ where $z \notin A$ and $B \subseteq A$ such that B and $A \setminus B$ are uncountable. Extend d to a metric d' on X by letting $d'(z, a) = 1$ if $a \in B$ and $d'(z, a) = 2$ if $a \in A \setminus B$. Since $B = \overline{B}(z, 1) \cap A$, B is a member of the relativized ball Borel structure, but is not a ball Borel set in (A, d) .

Despite this negative result, the relativization of the ball Borel structure of (X, d) to any bounded component X_0 is the ball Borel structure of (X_0, d) : This follows from the remark that any ball $B(x, r)$ with $r < \infty$, is either a subset of X_0 or is disjoint from X_0 .

There is no reason to expect pleasant behavior from the ball Borel structure for nonseparable metric spaces. However, the metric spaces of interest to us have the property that all bounded components are separable or equivalently, that all open balls of finite radius are separable. For instance, it follows immediately from Proposition 2.2, that all bounded components of a length space are σ -compact. In this case the relation between the ball Borel structure and the Borel structure is easily determined:

Proposition 2.3 *Suppose (X, d) is an extended metric space whose bounded components are all separable. Then the ball Borel structure of (X, d) is the σ -algebra $\mathcal{B}$ of those Borel sets A such that A or $\mathbb{C}A$ is a subset of a countable union of bounded components of X .*

PROOF. $\mathcal{B}$ is clearly a σ -algebra. Suppose A is a Borel set. If A is a subset of a bounded component X_0 of X , then by the equality of Borel structure and ball Borel structure on separable spaces, A is ball Borel in X_0 and therefore ball Borel in X . It follows that if A or $\mathbb{C}A$ is a subset of a countable union of bounded components of X , then A is ball Borel. Conversely, $\mathcal{B}$ contains all open balls of finite radius, so contains all ball Borel sets. ■

References for measure theory are ([7], [13]). We note one discrepancy between our notation and that used in these references. If μ is a countably additive measure on $(X, \mathcal{A})$ and $g : (X, \mathcal{A}) \rightarrow (Y, \mathcal{B})$ is measurable then we use the notation $g_*\mu$ to denote the measure μg^{-1} on $(Y, \mathcal{B})$.

2.2 Nonstandard Analysis

We need a very small amount of background material on nonstandard analysis such as the first few pages of Keisler's monograph [17]. Besides the Keisler reference, the book [1] is also highly recommended and will be referred to at

various places in the paper. Another general reference for this section with more foundational material is [15].

The *superstructure* over $\mathbf{G}$ is the set $V(\mathbf{G})$ defined by: $V_0(\mathbf{G}) = \mathbf{G}$, $V_{n+1}(\mathbf{G}) = V_n(\mathbf{G}) \cup \mathcal{P}(V_n(\mathbf{G}))$ and $V(\mathbf{G}) = \bigcup_n V_n(\mathbf{G})$. The main constituent of our working view of nonstandard analysis is a map $\star : V(\mathbb{R}) \longrightarrow V(\star \mathbb{R})$ which satisfies the transfer principle, c.f. [17] for details. In this paper we will only use the countable saturation property.

If $\mathfrak{X}$ is an internal set, $\text{card } \mathfrak{X}$ denotes the internal cardinality of $\mathfrak{X}$ if $\mathfrak{X}$ is hyperfinite, $+\infty$ otherwise.

If $r \in \star \mathbb{R}$, $r \ll \infty$ means that r is dominated by a standard real, $r \gg -\infty$ means that r dominates a standard real, r is *limited* iff $-\infty \ll r \ll \infty$, r is *infinitesimal* iff for every positive standard real θ , $|r| \leq \theta$. Hyperreals r, r' are *infinitely close*, written $r \cong r'$ iff $r - r'$ is infinitesimal. $r \ll r'$ means $r' - r$ is positive and not infinitesimal. The unique $r_0 \in \mathbb{R}$ infinitely close to r , if it exists is the *standard part* of r denoted $\text{st}(r)$. Define an (external) relation on $\star[0, \infty[$ by $r \preceq_O r'$ iff there is a limited hyperreal A such that $r \leq A r'$. Note that if $r' > 0$, then this is equivalent to $r/r' \ll \infty$. Similarly, define $r \preceq_o r'$ iff there is an infinitesimal hyperreal A such that $r \leq A r'$. Define $r \sim_O r'$ iff $r \preceq_O r'$ and $r' \preceq_O r$. The relation “ $\sim_O$ ” is clearly an equivalence relation, albeit an *external* one. If $r \sim_O r'$ we say r, r' are of the same order of magnitude. Note that for $r' > 0$, $r \sim_O r'$ iff $0 \ll r/r' \ll \infty$. Note also that $0 \sim_O r$ iff $r = 0$. If $r_i \sim_O r_1$, where $\{r_i\}_{1 \leq i \leq n}$, is a limited sequence of non-negative hyperreals, then $\sum_{i=1}^n r_i \sim_O r_1$.

Note that order of magnitude comparisons for individual real numbers are essentially vacuous. Similarly, for a function f defined on a finite interval of $\mathbb{R}$, one cannot usefully assign an order of polynomial growth to f . However, for functions on intervals in $\star \mathbb{R}$ the perspective of nonstandard analysis allows finer distinctions.

Definition 2.4 Suppose $0 < I_- \leq I_+$. A $\star \mathbb{R}$ -valued function f is of polynomial growth of order λ and scale factor σ in the interval $[I_-, I_+]$ iff $f(r) \sim_O \sigma r^\lambda$ for $r \in [I_-, I_+]$.

The scale factor and the order of growth are not uniquely determined. However:

Proposition 2.5 If f has polynomial growth on $[I_-, I_+]$ with $I_- \preceq_o I_+$, then the order of growth is uniquely determined up to $\cong$. If there is a hyperreal θ such that $0 \ll \theta \ll \infty$ and $I_- \leq (I_+/I_-)^\theta \leq I_+$, then the scale factor is unique up to $\sim_O$.

PROOF. Suppose $f(r) \sim_O \sigma r^\lambda \sim_O \sigma' r^\mu$ for $r \in [I_-, I_+]$. Assume without loss of generality that $\mu > \lambda$. Thus

$$\sigma/\sigma' \sim_O r^{\mu-\lambda} \tag{1}$$

for all $r \in [I_-, I_+]$ and so $I_+^{\mu-\lambda} \sim_O \sigma/\sigma' \sim_O I_-^{\mu-\lambda}$. Therefore,

$$\left(\frac{I_+}{I_-}\right)^{\mu-\lambda} \sim_O 1 \quad (2)$$

from which follows $(\mu - \lambda) \ln(I_+/I_-) \ll \infty$. Since $\ln(I_+/I_-) \cong \infty$, we conclude that $\mu - \lambda \cong 0$.

If $I_- \leq (I_+/I_-)^\theta \leq I_+$, then instantiating r in (1) with $(I_+/I_-)^\theta$, and using (2) and the fact $0 \ll \theta \ll \infty$, we deduce $\sigma/\sigma' \sim_O (I_+/I_-)^{\theta(\mu-\lambda)} \sim_O 1$. ■

2.3 Internal Metric Spaces

Suppose $(\mathfrak{X}, d)$ is an internal metric space. For $x, y \in \mathfrak{X}$, define $x \cong y$ iff $d(x, y)$ is infinitesimal. “ $\cong$ ” is an (external) equivalence relation on $\mathfrak{X}$ and $\diamond \mathfrak{X}$ is the quotient space $\mathfrak{X}/\cong$. For $x \in \mathfrak{X}$, let $\hat{x}$ be the $\cong$ -equivalence class of x in $\mathfrak{X}$. We will denote the canonical quotient map $x \rightarrow \hat{x}$ by $\varphi_{\mathfrak{X}}$ and refer to it as the *infinitesimal identification map*. The quotient space $\diamond \mathfrak{X}$ becomes an extended metric space with the extended metric $\diamond d(\hat{x}, \hat{y}) = \text{st}(d(x, y))$. $\diamond \mathfrak{X}$ with the metric $\diamond d$ is called the *infinitesimal hull* of $(\mathfrak{X}, d)$. In general $\diamond \mathfrak{X}$ is an extended metric space.

Example 2.1 If $N \cong \infty$, let $J = \{0, h, \dots, (N-1)h\}$, where $h = \frac{1}{N}$. The map $\text{st} : J \rightarrow [0, 1]$ is equivalent to the infinitesimal identification map, in the following sense: there is an isometric map $h : [0, 1] \rightarrow \diamond J$ such that $h \circ \text{st} = \varphi_J$.

We also define $x \sim y$ to mean $d(x, y) \ll \infty$. $\sim$ is also an external equivalence relation. The equivalence classes of $\mathfrak{X}$ under “ $\sim$ ” are the *limited components* of $\mathfrak{X}$. Clearly the bounded components of $\diamond \mathfrak{X}$ are the images under the map $\varphi_{\mathfrak{X}}$ of the limited components of $\mathfrak{X}$.

Example 2.2 Let $J' = \{-N h, -(N-1) h, \dots, 0, \dots, (N-1) h, N h\}$, where $h = \frac{1}{\sqrt{N}}$. In this case $\diamond J'$ is the disjoint union of its bounded components each one which is isometrically isomorphic to $\mathbb{R}$.

It follows immediately from countable saturation, that any separable metric space Y is isometrically isomorphic to a *subset* of a metric space $\diamond \mathfrak{X}$ with $\mathfrak{X}$ hyperfinite.

$\mathfrak{X} \mapsto \diamond \mathfrak{X}$ is a covariant functor from the category of internal metric spaces and S -continuous maps into the category of metric spaces and uniformly continuous mappings and $\varphi_{\mathfrak{X}} : \mathfrak{X} \rightarrow \diamond \mathfrak{X}$ is a natural transformation of functors.

We have the following relations between balls in $\mathfrak{X}$ and $\diamond \mathfrak{X}$: If $r' < r$ and both are standard, then

$$\overline{B}(x, r') \subseteq \varphi_{\mathfrak{X}}^{-1}(B(\hat{x}, r)) \subseteq B(x, r). \quad (3)$$

Similarly,

$$\overline{B}(x, r') \subseteq \varphi_{\mathfrak{X}}^{-1}(\overline{B}(\hat{x}, r')) \subseteq B(x, r). \quad (4)$$

For instance (4) follows from the implications $d(x, y) \leq r' \implies \text{st } d(x, y) \leq r' \implies d(x, y) < r$ which are valid so long as $r' < r$ are both standard.

The following proposition is well-known. See for instance [1].

Proposition 2.6 *If $(\mathfrak{X}, d)$ is an internal metric space, then each of the bounded components of $\diamond \mathfrak{X}$ is complete. In particular, $\diamond \mathfrak{X}$ is complete. If $C \subseteq \mathfrak{X}$ is internal, then $\diamond C$ (the image of C in $\diamond \mathfrak{X}$) is closed in $\diamond \mathfrak{X}$.*

For a metric space X in $V(\mathbb{R})$, define ι_X so that the following diagram commutes:

$$\begin{array}{ccc} X & & \\ \downarrow \star & \searrow \iota_X & \\ \star X & \xrightarrow{\varphi_{\star X}} & \diamond(\star X) \end{array}$$

Proposition 2.7 *If X is a $V(\mathbb{R})$ metric space, then $\iota_X : x \mapsto \varphi_{\star X}(\star x)$ is an isometric map $X \rightarrow \diamond(\star X)$. If X is compact, then ι_X is an isometric isomorphism $X \rightarrow \diamond(\star X)$.*

In particular, fix a (non-extended) metric space (X, d) . Since it is cumbersome to explicitly indicate the ι_X map $X \rightarrow \diamond(\star X)$, we usually view ι_X as an inclusion map with the property that $\varphi_{\star X}(\star x) = x$. Thus $\tilde{X} = \diamond(\star X)$ contains X itself and unless X is compact, $\tilde{X}$ is much larger than X . For example, consider $\mathbb{Z}$ with the metric $d(x, y) = 1$ for $x \neq y$. The ball $\overline{B}(0, 1) = \mathbb{Z}$ is not compact since the sequence $x_n = n$ has no convergent subsequences. Notice $\tilde{\mathbb{Z}}$ in this metric has one bounded component which is (much) larger than $\mathbb{Z}$. We now determine some conditions under which the bounded component of X in $\tilde{X}$ is X itself.

Proposition 2.8 *Suppose (X, d) is separable. A necessary and sufficient condition that the bounded component of X in $\tilde{X}$ coincide with X is that every closed ball of (X, d) be compact.*

PROOF. Observe first that if $a \in X$ is such that $\overline{B}(a, r)$ is compact, then $\overline{B}_{\tilde{X}}(a, r) = \overline{B}_X(a, r)$. Obviously, $\overline{B}_{\tilde{X}}(a, r) \supseteq \overline{B}_X(a, r)$. Let φ be the infinitesimal identification map $\star X \rightarrow \tilde{X}$. If $r < r'$, by formula (4) and well-known nonstandard characterizations of compactness ([1], 2.1.6),

$$\varphi^{-1}(\overline{B}_{\tilde{X}}(a, r)) \subseteq \overline{B}(\star a, r') = \star[\overline{B}(a, r')] \subseteq \varphi^{-1}(\overline{B}_X(a, r')).$$

Since φ is surjective, $\overline{B}_{\tilde{X}}(a, r) \subseteq \overline{B}_X(a, r')$, and as r' is an arbitrary standard real $> r$, $\overline{B}_{\tilde{X}}(a, r) \subseteq \overline{B}_X(a, r)$.

To show sufficiency, suppose $a \in X$ and let $x \in \tilde{X}$ be in the bounded component of a . There is a standard r such that $d_{\tilde{X}}(a, x) \leq r$. By the remarks of the first paragraph, $\overline{B}_{\tilde{X}}(a, r) = \overline{B}_X(a, r)$, so $x \in X$. Necessity: Suppose some ball $\overline{B}_X(a, r)$ with $r \in \mathbb{R}$ is not compact; let $\{y_i\}_{i \in \mathbb{N}}$ be a sequence of $\overline{B}_X(a, r)$

such that $\inf_{i \neq j} d(y_i, y_j) = \rho > 0$. $y_i = \hat{x}_i$ for some sequence $\{x_i\}_{i \in \mathbb{N}}$ of *X . By saturation this sequence extends to an internal sequence $x_1, \dots, x_N$ and by overspill we can also assume $d(x_i, x_j) \geq \rho/2$ for $1 \leq i, j \leq N$ with $i \neq j$ and $d({}^*a, x_i) \leq 2r$ for all i . In particular, $d(\hat{x}_i, \hat{x}_j) \geq \rho/2$ for $1 \leq i, j \leq N$ with $i \neq j$ and $d(a, \hat{x}_i) \leq 2r$ for all i . By assumption, the bounded component of X in $\bar{X}$ is X itself. Thus $\hat{x}_i \in X$ for all $1 \leq i \leq N$. Now $N \cong \infty$, so $\{1, \dots, N\}$ is uncountable. ■

Definition 2.9 *An internal mapping f from an internal metric space $(\mathfrak{X}, d_{\mathfrak{X}})$ to $(\mathcal{Y}, d_{\mathcal{Y}})$ is S -Lipschitz iff there is a limited hyperreal K such that for all $x, x' \in \mathfrak{X}$, $d_{\mathcal{Y}}(f(x), f(x')) \leq K d_{\mathfrak{X}}(x, x')$.*

There are related concepts such as S -Lipschitz equivalent metrics whose formulation we leave to the reader.

Finally, we note that the concept of length space is an internal one, so is applicable to standard metric spaces and internal metric spaces.

2.4 Borel Structure and Loeb Measure

We recall two basic concepts in nonstandard measure theory. The *inner Borel algebra* of an internal set $\mathfrak{X}$ is the smallest σ -algebra containing all internal subsets of $\mathfrak{X}$. A *hypermeasure* on $\mathfrak{X}$ is an internal, nonnegative and hyperfinitely additive function defined on all internal subsets of $\mathfrak{X}$. The conventional name given in the literature is the less mellifluous “hyperfinitely additive nonnegative measure”. The following result is essentially due to Loeb [19] in the case the hypermeasure is limited. The uniqueness result in the unlimited case is due to Henson (Corollary 1 of [14]).

Theorem 2.10 *Suppose μ is a hypermeasure on $\mathfrak{X}$. Then there is a unique countably additive measure on the inner Borel sets which extends the mapping $A \mapsto \text{st}(\mu(A))$ on the internal sets.*

The completion of the countably additive measure specified in the previous result is called the *Loeb measure* associated with μ . We denote Loeb measure by $L(\mu)$.

Proposition 2.11 *Suppose $(\mathfrak{X}, d)$ is an internal metric space. Then the infinitesimal identification map $\varphi_{\mathfrak{X}} : \mathfrak{X} \rightarrow \diamond \mathfrak{X}$ is measurable where $\mathfrak{X}$ has the inner Borel structure and $\diamond \mathfrak{X}$ has the ball Borel structure.*

PROOF. If $r \in \mathbb{R}$ then $\hat{y} \in B(\hat{x}, r) \iff \exists m \in \mathbb{N} \, d(x, y) < r - \frac{1}{m} \iff y \in \bigcup_{m \in \mathbb{N}} B(x, r - 1/m)$. Since each $B(x, r - 1/m)$ is internal, it follows $\varphi_{\mathfrak{X}}^{-1} B(\hat{x}, r)$ is inner Borel. ■

Definition 2.12 *Suppose $\mathfrak{X}$ is an internal metric space. If μ is a hypermeasure on $\mathfrak{X}$, then $\bar{P}(\mu)$ is the measure $\varphi_* L(\mu)$ defined on the σ -algebra of $A \subseteq \diamond \mathfrak{X}$ such that $\varphi^{-1}(A)$ is $L(\mu)$ measurable. $P(\mu)$ is the restriction of $\bar{P}(\mu)$ to the ball Borel sets of $\diamond \mathfrak{X}$.*

Note that since any inner Borel set is Loeb measurable, Propostion 2.11 immediately implies the ball Borel sets of $\diamond \mathfrak{X}$ are $\overline{P}(\mu)$ measurable.

Since $L(\mu)$ is complete, $\overline{P}(\mu)$ is also complete. Moreover, $\overline{P}(\mu)$ is not too far off from being the completion of $P(\mu)$.

Proposition 2.13 *Suppose μ is a hypermeasure on $\mathfrak{X}$ such that $P(\mu)(K) < \infty$ for every compact $K \subseteq \diamond \mathfrak{X}$. If $A \subseteq \diamond \mathfrak{X}$ is a subset of a σ -compact set and A is $\overline{P}(\mu)$ -measurable then A is in the completion of $P(\mu)$. Conversely, any set in the completion of $P(\mu)$ is $\overline{P}(\mu)$ measurable.*

PROOF. Suppose A is $\overline{P}(\mu)$ -measurable. By assumption, $A \subseteq \bigcup_n K_n$ with K_n compact. Since the domain of the completion of a measure is a σ -algebra, it suffices to show each $A \cap K_n$ is in the completion of $P(\mu)$. Thus without loss of generality, we can assume $A \subseteq K$ for some compact K . In particular,

$$L(\mu)(\varphi_{\mathfrak{X}}^{-1}(A)) = \overline{P}(\mu)(A) \leq P(\mu)(K) < \infty.$$

Let $\epsilon > 0$ with $\epsilon \in \mathbb{R}$ be arbitrary and $C \subseteq \varphi_{\mathfrak{X}}^{-1}(A)$ internal such that $L(\mu)(\varphi_{\mathfrak{X}}^{-1}(A) \setminus C) < \epsilon$. $\varphi_{\mathfrak{X}}(C)$ is closed and $\varphi_{\mathfrak{X}}(C) \subseteq A \subseteq K$ so $\varphi_{\mathfrak{X}}(C)$ is compact and hence ball Borel. Moreover,

$$\begin{aligned} \overline{P}(\mu)(A \setminus \varphi_{\mathfrak{X}}(C)) &= L(\mu)(\varphi_{\mathfrak{X}}^{-1}(A \setminus \varphi_{\mathfrak{X}}(C))) \\ &= L(\mu)(\varphi_{\mathfrak{X}}^{-1}(A) \setminus \varphi_{\mathfrak{X}}^{-1}(\varphi_{\mathfrak{X}}(C))) \\ &\leq L(\mu)(\varphi_{\mathfrak{X}}^{-1}(A) \setminus C) \leq \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, it follows there is a σ -compact (and therefore ball Borel set) $B \subseteq A$ such that $\overline{P}(\mu)(A \setminus B) = 0$. Applying this to $K \setminus A$, we conclude there is also a Borel set $B' \subseteq K$ and $\overline{P}(\mu)(B' \setminus A) = 0$. Thus A is nested between ball Borel sets B and B' with $\overline{P}(\mu)(B' \setminus B) = 0$, and so A is in the completion of $P(\mu)$. The converse follows from the definitions. ■

3 Compactness and Measure

We begin by giving some general compactness properties which follow from the existence of a Borel measure. All statements of this section are standard. If ν is a countably additive measure on a measurable space $(X, \mathcal{A})$, a set $A \subseteq X$ measurable or not is ν -finite iff there is a $B \in \mathcal{A}$ such that $A \subseteq B$ and $\nu(B) < \infty$. Any subset of a ν -finite set is ν -finite. In the context of the following result note [4].

Proposition 3.1 *Let (X, d) be a complete extended metric space and ν a countably additive ball Borel measure on X with the following property: There is an $R > 0$ such that*

$$\nu(B(x, r)) < \infty \quad \text{and} \quad 0 < \inf_{x, x'} \frac{\nu(B(x, r))}{\nu(B(x', r))}$$

for all $0 < r < R$. Then $K \subseteq X$ is relatively compact iff there is a $\delta > 0$ such that $\{x : d(x, K) \leq \delta\}$ is ν -finite.

PROOF. Suppose $K \subseteq X$ is relatively compact. Let $\epsilon \in \mathbb{R}$ be such that $0 < \epsilon < R$. By relative compactness of K , there is an $n \in \mathbb{N}$ and a sequence $x_1, \dots, x_n \in K$ such that $\overline{K} \subseteq \bigcup_{i=1}^n B(x_i, \epsilon) = W$. W is open, so $\delta = \inf\{d(x, y) : y \in X \setminus W, x \in \overline{K}\} > 0$. Let $V = \{x : d(x, K) \leq \delta/2\}$. Clearly $V \subseteq W$. Now $\bigcup_{i=1}^n B(x_i, \epsilon)$ is ball Borel and ν -finite. In particular V is ν -finite, proving the claim.

Conversely, suppose $\delta > 0$ is such that $V = \{x : d(x, K) \leq \delta\}$ is ν -finite. We may assume $\delta < R$. We prove by contradiction that K is precompact. By possibly choosing a smaller positive δ , we may assume there is an infinite sequence $\{x_i\}_{i \in \mathbb{N}}$ in K such that $x_i \notin B(x_j, \delta)$ for $j < i$. Thus $\{B(x_i, \delta/2)\}_{i \in \mathbb{N}}$ is a sequence of pairwise disjoint balls. Furthermore, $B(x_i, \delta/2) \subseteq V$, since $x_i \in K$. Thus for any N

$$\begin{aligned} \nu\left(\bigcup_{i=1}^{\infty} B(x_i, \delta/2)\right) &= \sum_{i=1}^{\infty} \nu(B(x_i, \delta/2)) \\ &\geq N \nu(B(x_1, \delta/2)) \inf_{1 \leq i \leq \infty} \frac{\nu(B(x_i, \delta/2))}{\nu(B(x_1, \delta/2))} \end{aligned}$$

so that $\nu\left(\bigcup_{i=1}^{\infty} B(x_i, \delta/2)\right) = +\infty$. Since $V \supseteq \bigcup_{i=1}^{\infty} B(x_i, \delta/2)$, V cannot be ν -finite. It follows that K is precompact. Thus by completeness of (X, d) the closure of K is compact. ■

Corollary 3.2 *Let (X, d) be a complete extended metric space and ν a ball Borel measure on X satisfying the conditions of Proposition 3.1. Then X is locally compact. In fact, all closed balls of radius $< R$ are compact.*

PROOF. Suppose $x \in X$ and r is a real with $0 < r < R$. Consider the set $V = \{y \in X : d(y, B(x, r)) \leq \delta\} \subseteq B(x, r + \delta)$. If $r + \delta < R$, then V is ν -finite. Thus by Proposition 3.1, $B(x, r)$ is relatively compact. ■

4 Near Homogeneity

Definition 4.1 *A hypermeasure μ on an internal metric space $(\mathfrak{X}, d)$ is nearly homogeneous iff there is an $R_\mu \gg 0$ such that for all open balls V, V' of radius s, s' respectively with $0 \ll s, s' \leq R_\mu$, $\mu(V) \sim_O \mu(V')$ and $\mu(V) \neq 0$. R_μ is a radius of homogeneity of μ .*

In this definition, note the possibility that R_μ is an unlimited hyperreal.

Denote the common $\sim_O$ -equivalence class of the hyperreals $\mu(B(x, s))$, for $0 \ll s \leq R_\mu$ by M_μ . Note that this is an external set. Elements of M_μ are normalization constants of μ . By abuse of notation, we write $r \sim_O M_\mu$ instead of $r \in M_\mu$. Similarly, we write $r \preceq_O M_\mu$ iff for any $s \in M_\mu$, $r \preceq_O s$. If $M_\mu \sim_O 1$, we say μ is *normalized*.

Given a nearly homogeneous hypermeasure μ on an internal metric space $(\mathfrak{X}, d)$, we associate to it a class of normalized hypermeasures on $\mathfrak{X}$ as follows:

For any $M \sim_O M_\mu$, let

$$\mu_M(A) = \frac{1}{M} \mu(A).$$

The hypermeasures μ_M are all essentially equivalent. For instance, the measures μ_M are constant multiples r , $0 \ll r \ll \infty$, of each other. It follows that the countably additive measures $L(\mu_M)$ have identical null sets and sets of finite measure.

We will call an internal metric space $(\mathfrak{X}, d)$ nearly homogeneous if the hypermeasure $\mu_{\text{card}} : A \mapsto \text{card } A$ is nearly homogeneous, where we make the convention that $\mu_{\text{card}}(A) = +\infty \in {}^*\mathbb{R}$ in case $A \subseteq \mathfrak{X}$ is not hyperfinite. Henceforth, if $(\mathfrak{X}, d)$ nearly homogeneous, we will use μ_{card} to denote a normalized version of $A \mapsto \text{card}(A)$.

In particular:

Proposition 4.2 *Suppose μ is a normalized nearly homogeneous hypermeasure on $(\mathfrak{X}, d)$. Then the measure $P(\mu)$ on $\diamond \mathfrak{X}$ has the property that*

$$0 < P(\mu)(B(x, r)) < \infty \quad (5)$$

and

$$0 < \inf_{x, x'} \frac{P(\mu)(B(x, r))}{P(\mu)(B(x', r))} \quad (6)$$

for all $r \in \mathbb{R}$ with $0 < r \leq R_\mu$.

PROOF. Let $\varphi : \mathfrak{X} \rightarrow \diamond \mathfrak{X}$ be the infinitesimal identification map. Thus

$$B(x, \frac{r}{2}) \subseteq \varphi^{-1}(B(\hat{x}, r)) \subseteq B(x, r).$$

Since μ is normalized, $\mu(B(x, \frac{r}{2})) \sim_O \mu(B(x, r)) \sim_O 1$, so

$$0 \ll \text{st } \mu(B(x, \frac{r}{2})) \leq P(\mu)(B(\hat{x}, r)) \leq \text{st } \mu(B(x, r)) \ll \infty.$$

This proves 5. To prove 6, note that

$$\frac{P(\mu)(B(\hat{x}, r))}{P(\mu)(B(\hat{x}', r))} \geq \text{st} \left(\frac{\mu(B(x, \frac{r}{2}))}{\mu(B(x', r))} \right).$$

However, for each $x, x' \in \mathfrak{X}$ and each standard r for which $0 < r \leq R_\mu$,

$$\frac{\mu(B(x, \frac{r}{2}))}{\mu(B(x', r))} \gg 0.$$

For a fixed value of r the above expression is an internal function of (x, x') . Therefore for a fixed standard value of r , its infimum over all pairs (x, x') is $\gg 0$ and (6) follows. ■

In particular:

Proposition 4.3 *Suppose μ is a normalized nearly homogeneous hypermeasure on $(\mathfrak{X}, d)$. All closed balls of radius $< R_\mu$ in $\diamond \mathfrak{X}$ are compact.*

5 Polynomial Growth

Definition 5.1 Let $(\mathfrak{X}, d)$ be an internal metric space, $I_- \leq I_+$ nonnegative hyperreals. $(\mathfrak{X}, d)$ is of polynomial growth in the interval $[I_-, I_+]$ iff there is a hyperreal σ and a hyperreal $0 \ll \lambda \ll \infty$ such that for $I_- \leq r \leq I_+$

$$\text{card } B(x, r) \sim_O \sigma r^\lambda. \quad (7)$$

The hyperreal σ is the *scale factor* and the limited hyperreal λ is the *order of growth* of $\mathfrak{X}$ in $[I_-, I_+]$. Alternatively, $(\mathfrak{X}, d)$ is of polynomial growth of order λ in $[I_-, I_+]$ with scale factor σ iff there are constants $c \gg 0$ and $C \ll \infty$ such that

$$c \sigma r^\lambda \leq \text{card } B(x, r) \leq C \sigma r^\lambda. \quad (8)$$

for $r \in [I_-, I_+]$. We will refer to the constants c, C as the lower and upper bounds of polynomial growth of $(\mathfrak{X}, d)$ in $[I_-, I_+]$.

Unless otherwise stated, we will implicitly assume $I_- \preceq_o I_+$. The following immediately follows from Proposition 2.5.

Proposition 5.2 If $I_- \preceq_o I_+$, then the order of growth is uniquely determined up to $\cong$. If there is a hyperreal θ such that $0 \ll \theta \ll \infty$ and $I_- \leq (I_+/I_-)^\theta \leq I_+$, the scale factor is unique up to $\sim_O$.

For each nonnegative $\gamma \in {}^*\mathbb{R}$, let $d_\gamma(x, y) = \frac{1}{\gamma}d(x, y)$. We use the notation B_γ to denote open balls relative to d_γ . Clearly, $d_\gamma(x, y) < r$ iff $d(x, y) < \gamma r$ so that $B_\gamma(x, r) = B(x, \gamma r)$.

Consider $\mathfrak{X}$ as a hypermeasure space with counting measure $\mu(V) = \text{card}(V)$.

Proposition 5.3 Suppose $I_- \preceq_o I_+$ and $\gamma > 0$ is such that $I_+/\gamma \gg 0$ and $\gamma/I_- \cong +\infty$. If $(\mathfrak{X}, d)$ is of polynomial growth in $[I_-, I_+]$ then counting measure μ is nearly homogeneous on $(\mathfrak{X}, d_\gamma)$ with a radius of homogeneity $R = I_+/\gamma$ and normalizing constant $M_\mu = \sigma \gamma^\lambda$.

In particular, closed balls of radius $< R$ are compact in $\diamond(\mathfrak{X}, d_\gamma)$ and $\diamond\mathfrak{X}$ is locally compact.

PROOF. If $0 \ll r, r' \leq R$, then $I_- \leq r\gamma, r'\gamma \leq R\gamma = I_+$, so

$$\text{card } B_\gamma(x, r) = \text{card } B(x, \gamma r) \sim_O \sigma (\gamma r)^\lambda \sim_O \sigma (\gamma r')^\lambda = \text{card } B_\gamma(x, r').$$

By Proposition 3.2, local compactness of $\diamond\mathfrak{X}$ follows. ■

If $(\mathfrak{X}, d)$ is an internal metric space of polynomial growth λ in $[I_-, I_+]$, then $\mu(A) = \text{card } A/\sigma \gamma^\lambda$ is a normalized homogeneous hypermeasure on $(\mathfrak{X}, d_\gamma)$. $\diamond(\mathfrak{X}, d_\gamma)$ is locally compact and is equipped with a countable additive ball Borel measure $P(\mu)$ which is finite on compacts.

Theorem 5.4 Suppose $(\mathfrak{X}, d)$ is of polynomial growth λ in $[I_-, I_+]$ and $\gamma > 0$ is such that $R = I_+/\gamma \gg 0$ and $\gamma/I_- \cong +\infty$. The countably additive measure $P(\mu)$ on $\diamond(\mathfrak{X}, d_\gamma)$ has the property that there are constants $k, K \in \mathbb{R}$ such that

$$k r^{\text{st}(\lambda)} \leq P(\mu)(B(x, r)) \leq K r^{\text{st}(\lambda)}$$

for every $x \in \diamond \mathfrak{X}$ and $r \in \mathbb{R}$ such that $0 < r \leq R$. Note that $B(x, r)$ refers to the ball in the metric $\diamond d_\gamma$.

PROOF. By definition, there are standard constants $0 < c \leq C < \infty$ such that for $0 \ll r \leq R$, $x \in \mathfrak{X}$

$$c \leq \frac{\text{card } B_\gamma(x, r)}{\sigma(r\gamma)^\lambda} \leq C$$

Now use formula (3) on $(\mathfrak{X}, d_\gamma)$. Thus if r is a nonnegative standard real such that $r \leq R$,

$$c \left(\frac{r}{2} \right)^{\text{st}(\lambda)} \leq \text{st } \mu(B_\gamma(x, \frac{r}{2})) \leq P(\mu)(B(\hat{x}, r)) \leq \text{st } \mu(B_\gamma(x, r)) \leq Cr^{\text{st}(\lambda)}.$$

Example 5.1 Consider a hyperfinite rooted tree T with leaf nodes $\text{leaf}(T)$. Any $x \in \text{leaf}(T)$ in T is uniquely identified by the sequence of branches $\text{path}(x) = \{b_i(x)\}_{1 \leq i \leq \text{depth}(x)}$ required to reach x from the root node of T . If m is a standard integer, define a metric d_m on $\text{leaf}(T)$ as follows: If $x \neq y$, $d_m(x, y) = m^{-v(x, y)}$ where $v(x, y)$ is the first index at which $\text{path}(x), \text{path}(y)$ differ, otherwise, $d_m(x, y) = 0$. It is well-known d_m is an ultrametric.

Proposition 5.5 *If T is a tree of uniform depth N all of whose non-leaf nodes have standard branching degree n , then for $r \in]m^{-(N+1)}, 1]$.*

$$n^{N-1} r^{\ln n / \ln m} \leq \text{card } B(x, r) \leq n^N r^{\ln n / \ln m}.$$

In particular, $(\text{leaf}(T), d_m)$ is of polynomial growth of order $\ln n / \ln m$ in the interval $]m^{-(N+1)}, 1]$.

PROOF. Note that the condition $r \in]m^{-(N+1)}, 1]$ is equivalent to

$$0 \leq -\ln r / \ln m < N + 1. \quad (9)$$

If $x \in \text{leaf}(T)$, then $B(x, r)$ consists of x and those $y \in \text{leaf}(T)$ with $x \neq y$ and $v(x, y) > -\ln r / \ln m$; $B(x, r)$ is thus the set of those y for which $b_i(x) = b_i(y)$ for $1 \leq i \leq -\ln r / \ln m$. Counting the number of all paths (from root to leaf node) which agree with $\text{path}(x)$ up to index $\lfloor -\ln r / \ln m \rfloor$ using (9), it follows that

$$\text{card } B(x, r) = n^{N - \lfloor -\ln r / \ln m \rfloor}.$$

The result now follows from the estimate

$$n^{N + \ln r / \ln m - 1} \leq \text{card } B(x, r) \leq n^{N + \ln r / \ln m},$$

and the fact that for any r , $n^{\ln r / \ln m} = r^{\ln n / \ln m}$. ■

Note that the Cantor middle thirds set has a Lipschitz equivalent leaf metric with $m = 3$ and $n = 2$. We also alert the reader to the fact that the spaces in this example are not length spaces, since they are ultrametric spaces and therefore totally disconnected.

6 Ahlfors Regularity

All results in this section are standard. $\mathcal{H}^\lambda$ denotes λ -dimensional Hausdorff measure, which is a countably additive measure on the Borel sets, and in particular on the ball Borel sets. For definition and background on Hausdorff measures, see [20], [23]. These references do not explicitly treat Hausdorff measures in extended metric spaces, but there is no difficulty extending the definitions and basic properties to this context. Nevertheless, we need to add the following caveat in case E is a subset of an extended metric space which does not have a countable δ cover (§1.2 of [8]). In this case, we make the convention that $\mathcal{H}_\delta^\lambda(E) = +\infty$ and accordingly, $\mathcal{H}^\lambda(E) = +\infty$.

Note that if $E \subseteq X$ is nonseparable, then $\mathcal{H}^\lambda(E) = +\infty$ for any λ . For if $\mathcal{H}^\lambda < \infty$ then for every $\delta > 0$ there is a countable set $D_\delta \subseteq X$ with the property that $d(x, D_\delta) \leq \delta$ for every $x \in E$. Thus $D = \bigcup_n D_{1/n}$ is countable and $d(x, D) = 0$ for all $x \in E$. In particular, an uncountable set which intersects each bounded component in exactly one point has λ -dimensional Hausdorff measure $+\infty$ for any λ . This oddity clearly suggests that we stick to separable subspaces whenever possible.

It is no great surprise that Theorem 5.4 determines the Hausdorff dimension of the extended metric space $\diamond(\mathfrak{X}, d_\gamma)$.

Proposition 6.1 *Suppose (X, d) is a complete extended metric space and ν a ball Borel measure on X . If there are real numbers $R > 0$ and $\lambda > 0$ such that for $0 < r < R$*

$$k r^\lambda \leq \nu(B(x, r)) \leq K r^\lambda \quad (10)$$

then all closed balls in X of radius $< R$ are compact. Moreover,

$$\nu(E) \leq K \mathcal{H}^\lambda(E) \quad (11)$$

for any ball Borel set E .

For the inequality in the other direction, there is a constant $c > 0$ such that for every Borel set E contained in a separable subset of (X, d) with $\nu(E) < \infty$, then $\mathcal{H}^\lambda(E) < \infty$ and

$$c \mathcal{H}^\lambda(E) \leq \nu(E) \quad (12)$$

PROOF. To show compactness of balls of radius $< R$, apply Corollary 3.1 to the complete metric space (X, d) and the ball Borel measure ν . We verify that the assumptions on ν hold: $\nu(B(x, r)) \leq K r^\lambda < \infty$ and for all $x, x' \in X$ and all $r < R$,

$$\frac{\nu(B(x, r))}{\nu(B(x', r))} \geq k/K > 0.$$

Let E be an arbitrary set, and $\{A_i\}_{i \in \mathbb{N}}$ any sequence which covers E . Then choosing $x_i \in A_i$,

$$\nu(E) \leq \sum_i \nu(A_i) \leq \sum_i \nu(B(x_i, \text{diam } A_i)) \leq K \sum_i \text{diam } A_i^\lambda.$$

Therefore, $\nu(E) \leq K \mathcal{H}_\delta^\lambda(E)$ for any positive δ and letting $\delta \rightarrow 0$, it follows that formula (11) holds for any ball Borel set E .

In order to show the inequality in the other direction, we need a polynomial upper bound on the sizes of coverings by balls. This is proved using well-known Vitali covering arguments. See for example [8], Theorem 1.10. For completeness (and since we have been unable to locate a reference with results in the precise form we need here,) we provide the details in the following subsection.

6.1 Vitali Covering Arguments

We begin with a lemma in which the cardinality bound on coverings is recast, by brute force, into a slightly more suggestive format:

Lemma 6.2 *Let (X, d) be a metric space, $a \in X$, $r > 0$ and $\lambda \in]0, \infty[$. Suppose there is a $\sigma_0 > 0$ such that for all $\sigma \leq \sigma_0$, there is an $Y_\sigma \subseteq X$ satisfying $d(z, Y_\sigma) < \sigma$ for all $z \in X$ and*

$$C = \sup_{\sigma \in]0, \sigma_0]} \frac{\text{card}(B(a, r) \cap Y_\sigma)}{\left(\frac{r}{\sigma}\right)^\lambda} < \infty. \quad (13)$$

Then the λ -dimensional Hausdorff measure of $B(a, r/2)$ is at most $2^\lambda C r^\lambda$.

PROOF. Suppose $\sigma \in]0, \min(\sigma_0, r/2)]$. Every $y \in B(a, r/2)$ is at distance $< \sigma$ from some $z_y \in Y_\sigma$; moreover by the triangle inequality and the fact $\sigma \leq r/2$, $z_y \in B(a, r)$. By formula (13) there are at most $C(r/\sigma)^\lambda$ distinct z_y . In particular, the ball $B(a, r/2)$ can be covered by the family $\{B(z_y, \sigma)\}_y$ of cardinality $\leq C(r/\sigma)^\lambda$. Each such ball has diameter $\leq 2\sigma$. Thus applying the definition of Hausdorff measure,

$$\mathcal{H}_\sigma^\lambda\left(B(a, \frac{r}{2})\right) \leq 2^\lambda \sigma^\lambda C \left(\frac{r}{\sigma}\right)^\lambda = 2^\lambda r^\lambda C.$$

This inequality is valid for $0 < \sigma \leq \min(\sigma_0, r/2)$. Thus the λ -dimensional Hausdorff measure of $B(a, r/2)$ is $\leq 2^\lambda r^\lambda C$ as claimed. ■

It remains to find a suitable supply of such sets Y_σ .

Definition 6.3 *An extended metric space (X, d) is σ -separated iff $d(x, y) \geq \sigma$ for all $x, y \in X$ with $x \neq y$.*

For any metric space (X, d) , by Zorn's lemma there is a set $Y \subseteq X$ which is maximal with respect to the property of being σ -separated. If $Y \subseteq X$ is maximal σ -separated, then for any $x \in X$, $d(x, Y) < \sigma$; otherwise, if $d(x, Y) \geq \sigma$, then $\{x\} \cup Y$ is also σ -separated.

Proposition 6.4 *Suppose (X, d) is a metric space, ν a ball Borel measure on X . Let $R > 0$ and $\lambda > 0$ be such that $\nu(B(a, R)) < \infty$ and $\nu(B(x, r)) \geq k r^\lambda$ for $x \in B(a, R/2)$ and every $r \leq R/2$. Then if $r \leq R/2$, the λ -dimensional Hausdorff measure of $B(a, r/2)$ is at most $4^\lambda k^{-1} \nu(B(a, 2r))$.*

PROOF. Let $r \leq R/2$. Suppose $0 < \sigma \leq r$ and Y_σ is maximal σ -separated. We will show

$$D = \frac{\text{card}(\mathcal{B}(a, r) \cap Y_\sigma)}{\left(\frac{r}{\sigma}\right)^\lambda} \leq 2^\lambda k^{-1} r^{-\lambda} \nu(\mathcal{B}(a, 2r)) < \infty,$$

independently of σ . The result will then follow by Lemma 6.2. Since Y_σ is σ -separated, the balls $\mathcal{B}(z, \sigma/2)$ for $z \in Y$ are pairwise disjoint. Moreover, $\mathcal{B}(a, 2r) \supseteq \bigcup_{z \in \mathcal{B}(a, r) \cap Y_\sigma} \mathcal{B}(z, \sigma/2)$. Thus

$$\nu(\mathcal{B}(a, 2r)) \geq \sum_{z \in \mathcal{B}(a, r) \cap Y} \nu(\mathcal{B}(z, \frac{\sigma}{2})) \geq D \left(\frac{r}{\sigma}\right)^\lambda k \left(\frac{\sigma}{2}\right)^\lambda = 2^{-\lambda} D k r^\lambda$$

The estimate on D follows. ■

Combining Lemmas 6.2 and 6.4 we very nearly have $\mathcal{H}^\lambda$ is absolutely continuous with respect to ν under the assumptions of Proposition 6.1. This is indeed true, as is shown by a Vitali covering argument. See for instance [8]. We state this as a lemma:

Lemma 6.5 *Let (X, d) be an extended metric space and $0 < \lambda < \infty$. Given an open set U in (X, d) and $\rho > 0$, define inductively a sequence of numbers $\{R_i\}_i$ and closed balls $V_i = \overline{\mathcal{B}}(x_i, r_i)$ with the following properties:*

1. $R_{m+1} = \sup\{r \leq \rho : \exists x \in X \text{ } \overline{\mathcal{B}}(x, r) \subseteq U \setminus \bigcup_{i=1}^m V_i\}$, where it is understood that if the set of real numbers in braces is empty then the sequences $\{R_i\}_i$ and $\{V_i\}_i$ terminate at m . Note that if R_{m+1} is defined, then it automatically follows that $R_{m+1} > 0$.
2. If $R_{m+1} > 0$, let x_{m+1}, r_{m+1} be such that $R_{m+1} \geq r_{m+1} > R_{m+1}/2$ and $\overline{\mathcal{B}}(x_{m+1}, r_{m+1}) \subseteq U \setminus \left(\bigcup_{i=1}^m V_i\right)$.

Then either $\sum_i r_i^\lambda = +\infty$ or $\mathcal{H}^\lambda(U - \bigcup_i V_i) = 0$.

PROOF. If the sequence $\{V_i\}_i$ is finite of length m , then $U = \bigcup_{i=1}^m V_i$. Suppose $\sum_i r_i^\lambda < \infty$. We show that for every $k \in \mathbb{N}$,

$$U = \bigcup_{i=1}^k V_i \cup \bigcup_{i=k+1}^{\infty} \overline{\mathcal{B}}(x_i, 3r_i). \quad (14)$$

To see this, suppose $x \in U \setminus \bigcup_{i=1}^k V_i$. U is open, so there is an $r > 0$ such that $\overline{\mathcal{B}}(x, r) \subseteq U \setminus \bigcup_{i=1}^k V_i$. Since $r_i \rightarrow 0$, there is an m such that such that $r_m < r/2$. $\overline{\mathcal{B}}(x, r)$ must intersect at least one of $V_{k+1}, \dots, V_{m-1}$. Otherwise $\overline{\mathcal{B}}(x, r) \subseteq U \setminus \bigcup_{i=1}^{m-1} V_i$ so that $R_m \geq r$ and hence $r_m > R_m/2 \geq r/2$ which contradicts $r_m < r/2$. Let ℓ be the first index $\geq k+1$ such that $\overline{\mathcal{B}}(x, r)$ intersects V_ℓ . Thus $R_\ell \geq r$ and $r_\ell > R_\ell/2 \geq r/2$. By the triangle inequality

$$d(x_\ell, x) \leq r_\ell + r \leq 3r_\ell,$$

so $x \in \bigcup_{i=k+1}^{\infty} \overline{B}(x_i, 3r_i)$ from which (14) follows. If $\delta > 0$ and k in (14) is such that $\text{diam } \overline{B}(x_i, 3r_i) = 6r_i < \delta$, then,

$$\mathcal{H}_{\delta}^{\lambda}(U \setminus \bigcup_{i=1}^{\infty} V_i) \leq 6^{\lambda} \sum_{i=k+1}^{\infty} r_i^{\lambda} \longrightarrow 0.$$

Thus $\mathcal{H}_{\delta}^{\lambda}(U \setminus \bigcup_{i=1}^{\infty} V_i) = 0$ and so taking the limit as $\delta \longrightarrow 0$, the result follows. ■

Completion of Proof of Proposition 6.1. We apply Lemma 6.4 as follows: If $r < R/2$, this lemma gives an upper bound on the λ -dimensional Hausdorff measure of any ball $B(x, r/2)$ for $r \leq R/2$ of $2^{3\lambda} k^{-1} K r^{\lambda}$.

Consider the case E is open with $\nu(E) < \infty$. Let $V_i \subseteq E$ be a disjoint sequence of closed balls of radius $r_i \leq R/2$ such that either $\sum_i r_i^{\lambda} = +\infty$ or $\mathcal{H}^{\lambda}(E \setminus \bigcup V_i) = 0$. In the first case,

$$\nu(E) \geq \sum_i \nu(V_i) \geq k \sum_i r_i^{\lambda} = +\infty,$$

which contradicts $\nu(E) < \infty$. Thus for some constants K_1, K_2, K_3 ,

$$\mathcal{H}^{\lambda}(E) = \sum_i \mathcal{H}^{\lambda}(V_i) \leq \sum_i K_1 r_i^{\lambda} \leq K_2 \sum_i \nu(V_i) \leq K_3 \nu(E).$$

Open balls of radius $\leq R/2$ have finite ν measure (by assumption) and finite $\mathcal{H}^{\lambda}$ measure by the preceding lemmas. By the first part of the theorem, balls of radius $\leq R/2$ are relatively compact, hence separable. By approximation properties of finite Borel measures (Theorem 1.1 of [16]), it follows that for open balls V of radius $\leq R/2$ inequality (12) holds for arbitrary Borel sets $E \subseteq V$. Since each separable Borel set E is a countable union of Borel sets contained in balls of radius $\leq R/2$, the general case follows. ■

Definition 6.6 *A complete extended metric space for which there are $0 < k \leq K < \infty$ and $R \in]0, \infty]$ such that*

$$k r^{\lambda} \leq \mathcal{H}^{\lambda}(B(x, r)) \leq K r^{\lambda} \quad (15)$$

for $0 < r < R$ is called Ahlfors regular of dimension λ up to R .

We emphasize that this definition allows $R = +\infty$. Note that Ahlfors regularity up to $+\infty$ is a property about the large scale structure of a space unlike its dimensional behavior which is a local property.

See [5] for more on this circle of ideas.

6.2 Discrete Characterization of Ahlfors Regularity

We can easily obtain a converse to Proposition 5.4.

Proposition 6.7 *A necessary condition a complete metric space (X, d) be Ahlfors regular of dimension λ up to some $R \in]0, \infty]$ is that*

1. *There exist an internal metric space $(\mathfrak{X}, d_{\mathfrak{X}})$ of polynomial growth λ on $[R_-, R_+]$ where $R_- \cong 0$ and $0 \ll R_+ \ll \infty$ if R is finite or R_+ is an unlimited hyperreal in case $R = \infty$*
2. *(X, d) is a subset of $\diamond(\mathfrak{X}, d_{\mathfrak{X}})$ with the property that for all $r < R$ and all $x \in X$, $B(x, r) \subseteq X$.*

Lemma 6.8 *Suppose μ satisfies inequality (10) for all $r \in]0, R[$. Then there are constants k', K' such that for all $\sigma > 0$ and $Y \subseteq X$ which is maximal σ -separated, the following holds:*

$$k' \leq \frac{\text{card}(B(a, r) \cap Y)}{\left(\frac{r}{\sigma}\right)^\lambda} \leq K' \quad \text{if } 2\sigma \leq r < R/2 \text{ and } a \in X. \quad (16)$$

PROOF. Let $D = D(a, r, \sigma, Y)$ be the expression in inequality (16) and let k and K be as in inequality (10). Since Y is σ -separated, the balls $B(z, \sigma/2)$ for $z \in Y$ are pairwise disjoint. Thus

$$\mu(B(a, 2r)) \geq \sum_{z \in B(a, r) \cap Y} \mu(B(z, \frac{\sigma}{2})) \geq D \left(\frac{r}{\sigma}\right)^\lambda k \left(\frac{\sigma}{2}\right)^\lambda = 2^{-\lambda} D k r^\lambda.$$

Since $2r < R$, inequality (10) and the previous inequality imply $D \leq 4^\lambda K k^{-1}$. In the other direction, $B(a, r/2) \subseteq \bigcup_{z \in B(a, r) \cap Y} B(z, \sigma)$. Thus,

$$\mu(B(a, \frac{r}{2})) \leq \sum_{z \in B(a, r) \cap Y} \mu(B(z, \sigma)) \leq D \left(\frac{r}{\sigma}\right)^\lambda K \sigma^\lambda.$$

Since $r/2 > 0$, inequality (8) again implies $D(a, r, \sigma, Y) \geq 2^{-\lambda} K^{-1} k$. ■

To complete the proof of Proposition 6.7, for each real number $\sigma > 0$, let Y_σ be a maximal σ -separated subset of X . Consider $*$ of the function $\sigma \rightarrow Y_\sigma$. By transfer, $[*Y]_\sigma \subseteq *X$ satisfies (16) and $[*Y]_\sigma$ is maximal σ separated in $*X$ for every positive hyperreal σ . Taking $\sigma \cong 0$, it follows $\diamond([*Y]_\sigma) = \diamond(*X)$. Now let $\mathfrak{X} = Y_\sigma$. $\mathfrak{X}$ satisfies (1) with $R_- = 2\sigma$ and $R_+ = R/2$. To prove (2), by Ahlfors regularity up to R of X , for $s < R$ the closed balls $\overline{B}(x, s)$ in X are compact and so $\overline{B}_{\diamond(*X)}(x, s) = \overline{B}(x, s) \subseteq X$. ■

The characterization of Ahlfors regular spaces given by Proposition 6.7 provides no information about the space $(\mathfrak{X}, d_{\mathfrak{X}})$. The remainder of the paper is devoted to obtaining a more informative result on the form of $(\mathfrak{X}, d_{\mathfrak{X}})$.

7 Graph Spaces

Graph means undirected graph, *without looping edges*. We denote the adjacency relation between vertices x, y in G by $x \longleftrightarrow y$. If G is connected, the *graph*

distance between $x, y \in G$ is the length of the shortest path in G from x to y . The graph distance is a metric with values in the nonnegative integers; we denote the graph distance by dist . For each $\gamma \in \mathbb{R}$, let $\text{dist}_\gamma(x, y) = \frac{1}{\gamma} \text{dist}(x, y)$. We use B to denote open balls relative to dist and B_γ to denote open balls relative to dist_γ .

We will be mainly considering the $\star$ -versions of graph-theoretic concepts. Vertices x, y of G belong to the same limited component of (G, dist_γ) iff there is a path from x to y of length $\preceq_O \gamma$. In particular, if the graph diameter of (G, dist_γ) is $\preceq_O \gamma$, then (G, dist_γ) has exactly one limited component.

We contrast our notion of polynomial growth in metric spaces with the standard one for graphs: If G is a graph, G has standard polynomial growth $\lambda \in \mathbb{R}$ iff there are constants $0 < c \leq C < \infty$ in $\mathbb{R}$ such that

$$c \ell^\lambda \leq \text{card} B(x, \ell) \leq C \ell^\lambda \quad \text{for } \ell \geq 1. \quad (17)$$

Note that standard polynomial growth for internal graphs is an external property.

Example 7.1 There is a vast literature on growth of finitely generated discrete groups leading to Gromov's theorem on virtually nilpotent groups (See [2], [10], [9], [24].) See [12] for definitions and examples of nilpotent groups.

We will need a basic result on nilpotent groups due to Bass [2]. Recall first that for a group G , and $V \subseteq G$ such that $V = V^{-1}$ and $e \notin V$, the *Cayley graph* of G relative to V , denoted $\text{Cayley}(G, V)$, is the undirected loopless graph whose vertices are the elements of G and whose edges are the pairs $\{x, y\}$ such that $x^{-1}y \in V$. If G is a finitely generated nilpotent group, then for any symmetric generating set W , there exists a $d(G) \in \mathbb{N}$ such that $\text{Cayley}(G, W)$ has standard growth $d(G)$. Theorem 2 of [2], provides an explicit formula for $d(G)$.

Proposition 7.1 *Suppose G is an internal nilpotent group of class $n \in \mathbb{N}$. Suppose furthermore $W \subseteq G$ is a symmetric set with $\text{card} W \in \mathbb{N}$ and such that W generates G as an internal group. Then there is a standard integer λ such that $\text{Cayley}(G, W)$ is of polynomial growth λ on some interval $[2, M]$ for $M \cong \infty$.*

PROOF. The external group G_∞ generated by W is nilpotent of class $\leq n$ and finitely generated. By Bass's result, $\text{Cayley}(G_\infty, W)$ is of standard polynomial growth $\lambda \in \mathbb{N}$. This means that for some $c, C \in]0, \infty[$,

$$c \ell^\lambda \leq \text{card}(B(e, \ell) \cap G_\infty) \leq C \ell^\lambda \quad (18)$$

for all $\ell \in \mathbb{N}$. For $\ell \in \mathbb{N}$, $B(e, \ell) \cap G_\infty = B(e, \ell)$. Thus (18) actually implies an internal condition, i.e. inequality (17) with $x = e$. Thus by overspill there is an $R_+ \cong \infty$ such that (17) holds with $x = e$ for all $\ell \leq R_+$. However, $\text{card} B(e, \ell) = \text{card} B(x, \ell)$ for any $x \in G$, so $\text{Cayley}(G, W)$ is of polynomial growth λ on some interval $[2, M]$ for $M \cong \infty$. ■

8 Lifting Measures

Proposition 8.1 *Let (X, d_X) be a σ -compact metric space, μ a ball Borel measure on X such that $\mu(K) < \infty$ for every compact $K \subseteq X$, $(\mathfrak{X}, d_{\mathfrak{X}})$ a hyperfinite space such that $\diamond \mathfrak{X} \supseteq X$. Then there is a hypermeasure ν on $\mathfrak{X}$ such that $P_{\mathfrak{X}}(\nu) \upharpoonright \diamond \mathfrak{X} = \mu$.*

PROOF. This is a special case of the discussion in [1], §5.2 on lifting measures. The arguments used there rely on more saturation than we need in this paper, but given the separability assumptions on (X, d_X) we can get by with only countable saturation. For completeness we give a direct proof assuming only countable saturation.

Assume first X is compact. We may assume without loss of generality that μ is a probability measure and $\mathfrak{X} = {}^*X$. To show this last remark, note there is a hyperfinite $F \subseteq \mathfrak{X}$ such that $\diamond F = X$ and a mapping $g : {}^*X \rightarrow F$ which makes the following diagram commutative:

$$\begin{array}{ccc} {}^*X & \xrightarrow{g} & F \\ \varphi_{*X} \downarrow & & \downarrow \varphi_F \\ X & \xrightarrow{\text{id}_X} & X \end{array}$$

Thus any lifting for μ in *X can be pushed over by g to a lifting for μ in F . There is a countable set $\mathcal{K}$ of compact sets in X such that every open set in X is the union of a nondecreasing sequence of $\mathcal{K}$. By the saturation property, there is a hyperfinite boolean algebra $\mathcal{F} \subseteq {}^*\mathfrak{B}$, where $\mathfrak{B}$ is the Borel algebra of the compact metric space X , such that for each $K \in \mathcal{K}$, ${}^*K \in \mathcal{F}$. Now there is an internal operator κ on the limited nonnegative hyperfinitely additive set functions on $\mathcal{F}$ such that for any limited hyperfinitely additive set function ν , $\kappa(\nu)$ extends ν and is a limited hypermeasure on *X . ${}^*\mu$ is a limited nonnegative hyperfinitely additive set function on the algebra ${}^*\mathfrak{B}$. Let $\nu = \kappa({}^*\mu \upharpoonright \mathcal{F})$ and $\rho = P(\nu)$. ${}^*\mu$ has total mass one and thus ν has total mass one. ρ is a Borel measure on X of total mass ≤ 1 . For any $K \in \mathcal{K}$, ${}^*K \subseteq \varphi^{-1}(K)$ so that

$$\rho(K) = L(\nu)(\varphi^{-1}(K)) \geq L(\nu)({}^*K) = \text{st}({}^*\mu({}^*K)) = \mu(K).$$

By monotonicity of measures, and the fact every open set is the union of a nondecreasing sequence of $\mathcal{K}$, it follows that $\rho(U) \geq \mu(U)$ for any open set U . By regularity (see [16], Theorem 1.4), $\rho(E) \geq \mu(E)$ for any Borel set E . Since μ is a probability measure and $\rho(X) \leq 1$, ρ is also a probability measure. It follows $\rho = \mu$.

In the general case, there is a nondecreasing sequence of compact sets $\{K_i\}_{i \in \mathbb{N}}$ such that $X = \bigcup_i K_i$. For each $i \in \mathbb{N}$, let $E_i \subseteq \mathfrak{X}$ be hyperfinite and such that $\diamond E_i \supseteq K_i$. Existence of the sets E_i follows immediately from countable saturation. Clearly we may assume $E_i \subseteq E_{i+1}$. By assumption, $\mu \upharpoonright K_i$ is a finite Borel measure. By the special case considered in the previous paragraph, there is a sequence of limited hypermeasures $\{\nu_i\}_{i \in \mathbb{N}}$ such that ν_i is supported in E_i

and $P(\nu_i) = \mu|K_i$. We may assume $\nu_i \leq \nu_{i+1}$. Proof: Consider $\mu|K_i$ as a measure on K_{i+1} by assigning total measure 0 to $K_{i+1} \setminus K_i$. With this convention, $\mu|K_i \leq \mu|K_{i+1}$, so that $\mu|K_{i+1} - \mu|K_i$ is also a finite countably additive measure. Now let ν'_{i+1} be such that $P(\nu'_i) = \mu|K_{i+1} - \mu|K_i$ and ν'_{i+1} is supported in E_{i+1} . Then $\nu_{i+1} = \nu'_{i+1} + \nu_i$ is the desired measure.

$\varphi^{-1}(K_i)$ is Borel, so by approximation of Loeb measurable sets of finite measure by internal subsets, there is an internal set F_i such that $\varphi^{-1}(K_i) \subseteq F_i$ and

$$\nu_{i+1}(F_i) \leq L(\nu_{i+1})(\varphi^{-1}(K_i)) + 2^{-(i+1)}.$$

Now

$$L(\nu_{i+1})(\varphi^{-1}(K_i)) = P(\nu_{i+1})(K_i) = \mu(K_i) = P(\nu_i)(K_i) = L(\nu_i)(\varphi^{-1}(K_i)),$$

so

$$\nu_{i+1}(F_i) \leq L(\nu_i)(\varphi^{-1}(K_i)) + 2^{-(i+1)} \leq L(\nu_i)(F_i) + 2^{-(i+1)} \leq \nu_i(F_i) + 2^{-i}.$$

Since $\nu_i \leq \nu_{i+1}$ it follows that for all internal $A \subseteq F_i$,

$$\nu_i(A) \leq \nu_{i+1}(A) \leq \nu_i(A) + 2^{-i}.$$

Now apply saturation and overspill, to conclude that the sequences ν_k , E_k and F_k have extensions to hyperfinite sequences defined for $1 \leq k \leq N$ such that

1. $E_i \subseteq E_{i+1}$ for $i \leq N-1$,
2. E_i is hyperfinite and $E_i \subseteq F_i$ for $i \leq N$,
3. $\nu_i(A) \leq \nu_{i+1}(A) \leq \nu_i(A) + 2^{-i}$ for $i \leq N-1$ and all internal $A \subseteq F_i$.

It follows from item 3, that for $j \geq i$,

$$\nu_i(A) \leq \nu_j(A) \leq \nu_i(A) + 2^{-(i-1)} \tag{19}$$

for all internal $A \subseteq F_i$. It follows immediately from (19) and the monotone class theorem that for any inner Borel set $A \subseteq F_i$,

$$L(\nu_i)(A) \leq L(\nu_j)(A) \leq L(\nu_i)(A) + 2^{-(i-1)}.$$

Let $\nu = \nu_N$. We need to show that for any Borel set $A \subseteq \diamond \mathfrak{X}$, $P(\nu)(A) = \mu(A)$. If $i \in \mathbb{N}$, then for any ball Borel set $A \subseteq K_i$, $\varphi^{-1}(A) \subseteq \varphi^{-1}(K_i) \subseteq F_i$, so

$$\begin{aligned} \mu(A) = P(\nu_i)(A) &= L(\nu_i)(\varphi^{-1}(A)) \\ &\leq L(\nu)(\varphi^{-1}(A)) \\ &= P(\nu)(A) \\ &\leq L(\nu_i)(\varphi^{-1}(A)) + 2^{-(i-1)} \\ &= P(\nu_i)(A) + 2^{-(i-1)} = \mu(A) + 2^{-(i-1)} \end{aligned}$$

Since i is arbitrary, $\mu(A) = P(\nu)(A)$ for any Borel set A which is a subset of some K_i . However, $\bigcup_{i \in \mathbb{N}} K_i = X$, so by countable additivity of μ and of $P(\nu)$, the result follows for all Borel sets $A \subseteq X$. ■

A hypermeasure on the internal subsets of $\mathfrak{X}$ is *uniform* if all the singleton sets have the same mass. Define the *density* of ν to be the mass of each singleton.

Proposition 8.2 *Let ν be a hypermeasure on a hyperfinite set $\mathfrak{X}$. Then there is a uniform hypermeasure μ on a hyperfinite set $\mathcal{Y}$ and an internal mapping $p : \mathcal{Y} \rightarrow \mathfrak{X}$ such that $\mu(p^{-1}(A)) \cong \nu(A)$ for every internal $A \subseteq \mathfrak{X}$ and $\mu(\mathcal{Y}) \leq \nu(\mathfrak{X})$.*

PROOF. Assume $\mathfrak{X} = \{1, \dots, N\}$ and let $\nu_1, \dots, \nu_N$ be the masses of the atoms of $\mathfrak{X}$. Let M be such that $M/N \cong \infty$. For each $i \in \{1, \dots, N\}$ there is a unique hyperinteger $0 \leq k_i$ such that $k_i/M \leq \nu_i < (k_i + 1)/M$. Let $\{B_i\}_{1 \leq i \leq N}$ be disjoint sets such that $\text{card}(B_i) = k_i$, and define $p : \mathcal{Y} \rightarrow \mathfrak{X}$ so that p is identically i on B_i , for $i \geq 1$. Let μ be the hypermeasure on $\mathcal{Y}$ which assigns the mass $1/M$ to each singleton of $\mathcal{Y}$. For every internal $A \subseteq \mathfrak{X}$,

$$\mu(p^{-1}(A)) = \sum_{i \in A} \mu(B_i) = \sum_{i \in A} k_i/M = \sum_{i \in A} (\nu_i + o_i) = \nu(A) + \sum_{i \in A} o_i$$

Now $|\sum_{i \in A} o_i| \leq \sum_{i \in \mathfrak{X}} |o_i| \leq N/M \cong 0$. Finally $\mu(\mathcal{Y}) = \sum_i k_i/M \leq \sum_i \nu_i = \nu(\mathfrak{X})$. ■

It is clear that there is a lot of leeway in the choice of density $1/M$. In particular, M can be chosen to be a hyperinteger. In case μ is a hypermeasure on $\mathfrak{X}$ and $(\mathfrak{X}, d)$ is an internal metric space, we can take $\mathcal{Y}$ to be an internal metric space. In case $(\mathfrak{X}, d)$ is a graph metric space we can $\mathcal{Y}$ to be a graph metric space also.

Definition 8.3 *A spiked graph over G is a family of connected loopless graphs $\{H_v\}_{v \in \text{nodes}(G)}$ which are pairwise disjoint and such that $v \in H_v$ for each node v of G .*

The disjoint union of a spiked graph is the loopless graph G' obtained as follows: The set of nodes of G' is the disjoint union $\bigsqcup_v \text{nodes}(H_v)$. An edge of G' is either an edge in G or an edge in one of the graphs H_v . We will use the expression $\bigsqcup_v H_v$ to denote the disjoint union.

The base projection of the disjoint union is the mapping which is identically v on H_v .

We will use informal but suggestive terminology to describe spiked graphs: The graphs H_v are the spikes, etc.. Note that the disjoint union of a spiked graph over G contains G as a full subgraph. A *spike extension* of a graph G is a loopless graph G' which is the disjoint union of a spiked graph over G .

Proposition 8.4 *Under the assumptions of Proposition 8.2, if $(\mathfrak{X}, d_{\mathfrak{X}})$ is an internal metric space, we may assume $(\mathcal{Y}, d_{\mathcal{Y}})$ is an internal metric space and $p : \mathcal{Y} \rightarrow \mathfrak{X}$ satisfies*

$$d_{\mathfrak{X}}(p(y), p(y')) \leq d_{\mathcal{Y}}(y, y') \leq d_{\mathfrak{X}}(p(y), p(y')) + \epsilon(y, y')$$

where $\epsilon(y, y') \cong 0$.

Suppose $d_{\mathfrak{X}}$ is $\delta \times \text{dist}$ where $\delta \cong 0$ and dist the metric associated to a loopless graph on $\mathfrak{X}$. Then $d_{\mathcal{Y}}$ can also be taken as $\delta \text{dist}'$ for some metric associated to a loopless graph on $\mathcal{Y}$ which is a spike extension of $\mathfrak{X}$ and p is the base projection.

PROOF. Provide each set $p^{-1}(x)$ with a metric d'_x which assumes only infinitesimal values and with a distinguished point y_x . The metric d'_x can be chosen in a completely arbitrary way, but one possible choice is as follows: Choose an infinitesimal θ and $d'_x(a, b) = \theta$, whenever $a \neq b$. Define

$$d_{\mathcal{Y}}(y, y') = d_{\mathfrak{X}}(p(y), p(y')) + d'_{p(y)}(y, y_{p(y)}) + d'_{p(y')}(y_{p(y')}, y')$$

It is clear $d_{\mathcal{Y}}$ has the desired properties.

In the case $d_{\mathfrak{X}} = \delta \times \text{graph metric}$ on $\mathfrak{X}$, consider $\mathcal{Y}$ as a loopless graph obtained by attaching spikes to $\mathfrak{X}$, where now each spike is a complete graph. Then $p : \mathcal{Y} \rightarrow \mathfrak{X}$ is the mapping which maps each member of $\mathcal{Y}$ to the attachment point of the spike in $\mathfrak{X}$. Since each spike is a complete graph, $\text{dist}(y, y') \leq 1$ for all y, y' on a spike, the metric $d_{\mathcal{Y}}$ on each spike is infinitesimal. ■

The space $\mathcal{Y}$ of the preceding result can be regarded as an infinitesimal thickening of $\mathfrak{X}$. Note that also the following diagram commutes

$$\begin{array}{ccc} \mathcal{Y} & \xrightarrow{p} & \mathfrak{X} \\ \varphi_{\mathcal{Y}} \downarrow & & \downarrow \varphi_{\mathfrak{X}} \\ \diamond \mathcal{Y} & \longrightarrow & \diamond \mathfrak{X} \end{array}$$

where the bottom arrow is an isometric isomorphism.

8.1 Graph Regularity

Nothing has been said about a *regular* graph structure on $\mathcal{Y}$ (e.g., all nodes have the same degree), but this can also be arranged.


Proposition 8.5 *Under the assumptions of Proposition 8.2, suppose $d_{\mathfrak{X}}$ is $\delta \times \text{dist}$ where $\delta \cong 0$ and dist is the metric associated to a loopless graph on $\mathfrak{X}$. Then $d_{\mathcal{Y}}$ can also be taken as $\delta \text{dist}'$ for some metric associated to a regular loopless graph on $\mathcal{Y}$ which is a spike extension of $\mathfrak{X}$.*

PROOF. By Proposition 8.4, we may assume ν is a uniform hypermeasure on $\mathfrak{X}$ with density M . Let $k \in {}^*\mathbb{N}$ be odd and such that $k \geq 3 + \max_{x \in \mathfrak{X}} \deg_{\mathfrak{X}}(x)$. Since $\mathfrak{X}$ is hyperfinite, such a k exists. There exists a spiked graph over $\mathfrak{X}$, $\{F_x\}_{x \in \mathfrak{X}}$ such that

1. $\text{card } F_x = k + 3$,
2. For each $x \in \mathfrak{X}$, x has degree $k + 1 - \deg_{\mathfrak{X}}(x)$ in F_x ,
3. For each $x \in \mathfrak{X}$ and all $y \in F_x \setminus \{x\}$, y has degree $k + 1$ in F_x .

Then the loopless graph $\mathcal{Y} = \bigsqcup_{x \in \mathfrak{X}} F_x$ (see Definition 8.3) is regular of degree $k + 1$. If dist' is the graph distance on $\mathcal{Y}$, then each $x \in \mathfrak{X}$ is at dist' distance at most 2 from every $y \in F_x$. To see this, note that by the degree assumption on x , x is connected to at least one $y_x \in F_x \setminus \{x\}$. By the degree assumption on y , every $y \in F_x \setminus \{x\}$ is connected to all other $y' \in F_x$ but one. If y is connected to x , we are done. Otherwise, y is connected to y_x . It follows that each $x \in \mathfrak{X}$ is at infinitesimal $d_{\mathcal{Y}}$ distance from every $y \in F_x$. Let ν' be the uniform hypermeasure on $\mathcal{Y}$ with density $M/(k + 3)$.

We need to prove that for each x there is a loopless graph F_x with the required property. Start off with a disjoint family of sets $\{F_x\}_{x \in \mathfrak{X}}$ such that $x \in F_x$ and $\text{card } F_x = k + 3$. We will add edges to F_x so that the above requirements are met. Partition F_x into the sets $\{x\}$, A and B as shown in the figure, where $s = k + 1 - \deg_{\mathfrak{X}}(x) \geq 4$:



The line on the right represents a set of s edges joining x to each of the s members of A . As the following lemma shows, we can add edges between members of A so that each element of A is joined to exactly $s - 2$ other elements of A :

Lemma 8.6 *Suppose $\text{card } A \geq 4$ is even and $r \leq \text{card } A - 2$ is even. Then there is a connected loopless graph on A which is regular of degree r .*

PROOF. Let $s = \text{card } A$ and consider the additive group $\mathbb{Z}/(s)$. $\mathbb{Z}/(s)$ is a cyclic group of order s . Let V be the subset of $\mathbb{Z}/(s)$ consisting of the equivalence classes of $\pm 1, \dots, \pm r/2$. V has cardinality r . The Cayley graph $\text{Cayley}(\mathbb{Z}/(s), V)$ is regular of degree r . Since $0 \notin V$, $\text{Cayley}(\mathbb{Z}/(s), V)$ is also loopless. ■

Each element of A so far has attached $s - 1$ edges : $s - 2$ connected to other members of A and 1 edge to x . To get up to $k + 1$ edges per node, we must leave $k + 1 - (s - 1) = k + 2 - s$ unattached out edges for each of the s members of A . This is a total of $s(k + 2 - s)$ dangling edges (to the left) from A . Similarly each element of B can be joined to $k + 1 - s$ nodes of B — just consider the complete graph on B — leaving $k + 1 - (k + 1 - s) = s$ unattached out edges per element of B . This is a total of $s(k + 2 - s)$ dangling edges (to the right) from B . Now attach each right edge from B to a unique left edge from A . ■

9 Spaces from Graphs

Example 9.1 Consider $\mathbb{R}^n$ with the metric $d(\vec{x}, \vec{y}) = \sum_k |x_i - y_i|$. This metric is sometimes called the Taxicab distance [18]. Closed n -cubes $\prod_{i=1}^n [a_i, b_i]$ in $\mathbb{R}^n$ have the length space property with respect to the Taxicab metric.

Example 9.2 More generally, in the case $n \geq 2$ it is geometrically clear that the result of removing a finite number of open n -cubes from a closed n -cube is also a length space. This example easily extends to the removal of a countable number of disjoint open n -cubes.

Example 9.3 Suppose G is a hyperfinite loopless graph with graph distance dist . For any $M \cong \infty$, $\diamond(G, \text{dist}_M)$ is a length space. Proof: Suppose $\hat{x}, \hat{y} \in \diamond G$ are such that $\text{st}(\text{dist}_M(x, y)) = \diamond \text{dist}_M(\hat{x}, \hat{y}) = a$. Thus there is a path $x = z_0 \longleftrightarrow z_1 \cdots \longleftrightarrow z_n = y$ with $n = \text{dist}(x, y)$ and $a = \text{st}(n/M)$. The map $T = \{0, 1/M, \dots, n/M\} \rightarrow (G, \text{dist}_M)$ given by $j/M \mapsto z_j$ is clearly isometric and thus factors through an isometry $[0, a] \rightarrow \diamond(G, \text{dist}_M)$.

In this section we will characterize Ahlfors regular length spaces (X, d_X) as bounded components of hulls of hyperfinite spaces $(\mathfrak{X}, d)$. The main technical point of the proof is to show that a certain subset $\mathcal{Y}$ of $\mathfrak{X}$ has polynomial growth. Specifically, we need estimates of the kind

$$k r^\lambda \leq \text{card}(\text{B}(x, r) \cap \mathcal{Y}) \leq K r^\lambda$$

for $x \in \mathcal{Y}$. Note that this lower bound is stronger than the lower bound $k r^\lambda \leq \text{card} \text{B}(x, r)$.

Lemma 9.1 *Suppose (X, d) is a length space and $\rho > 0$. Let $a, x \in X$ be such that $d(a, x) \leq \rho$. Then for all r such that $0 \leq r \leq \rho$, there is a $c \in X$ such that*

$$\text{B}(c, \frac{r}{2}) \subseteq \text{B}(a, \rho) \cap \text{B}(x, r). \quad (20)$$

PROOF. We consider two cases:

Suppose $d(a, x) < r/2$. In this case take $c = x$. Obviously, $\text{B}(x, r/2) \subseteq \text{B}(x, r)$. If $y \in \text{B}(x, r/2)$, then $d(a, y) \leq d(a, x) + d(x, y) < r/2 + r/2 = r \leq \rho$ so that $\text{B}(x, r/2) \subseteq \text{B}(a, \rho)$.

Suppose $d(a, x) \geq r/2$. In this case we need the length space property of (X, d) . Let $\rho' = d(a, x)$. There is an isometric map $f : [0, \rho'] \rightarrow X$ with $f(0) = a$ and $f(\rho') = x$. Since $0 \leq \rho' - r/2 \leq \rho'$, $c = f(\rho' - r/2)$ is well-defined. Now $d(a, c) = \rho' - r/2$ and $d(c, x) = |\rho' - r/2 - \rho'| = r/2$. Thus by the triangle inequality, $\text{B}(c, r/2) \subseteq \text{B}(x, r)$ and $\text{B}(c, r/2) \subseteq \text{B}(a, \rho') \subseteq \text{B}(a, \rho)$. ■

Lemma 9.2 *Suppose (X, d) is an Ahlfors regular length space of dimension λ up to R . Then there exists $k' > 0$ such for every $a, x \in X$ and $r, \rho > 0$ satisfying $r < R$ and $r \leq \rho$ and $d(x, a) \leq \rho$,*

$$k' r^\lambda \leq \mathcal{H}^\lambda(\text{B}(x, r) \cap \text{B}(a, \rho))$$

PROOF. Let k be as in Formula (15). By the preceding lemma there is a c satisfying the inclusion (20). Thus,

$$k \left(\frac{r}{2}\right)^\lambda \leq \mathcal{H}^\lambda(\text{B}(c, \frac{r}{2})) \leq \mathcal{H}^\lambda(\text{B}(x, r) \cap \text{B}(a, \rho))$$

so $k' = 2^{-\lambda} k$ will do. ■

Proposition 9.3 *Suppose (X, d_X) is an Ahlfors regular length space of dimension λ up to $R \in]0, \infty]$, $(\mathfrak{X}, d_{\mathfrak{X}})$ is a hyperfinite space such that*

1. *(X, d_X) is a bounded component of $\diamond(\mathfrak{X}, d_{\mathfrak{X}})$,*
2. *For some uniform hypermeasure ν on $\mathfrak{X}$, $P_{\mathfrak{X}}(\nu)|_X = \mathcal{H}^\lambda$.*

Then there is an internal $\mathcal{Y} \subseteq \mathfrak{X}$ such that $\diamond \mathcal{Y} \supseteq X$ and $(\mathcal{Y}, d_{\mathfrak{X}}|_{\mathcal{Y}})$ is of polynomial growth λ on some interval $[R_-, R]$ with $R_- \cong 0$ in case R is finite or $[R_-, R_\infty]$ for some unlimited R_∞ and $R_- \cong 0$ in case $R = \infty$.

PROOF. Choose $a \in \mathfrak{X}$ so that $\hat{a} \in X$. In particular, X is the bounded component of $\hat{a}$. For any standard r , standard ρ and $x \in \mathfrak{X}$ the Formula (3) implies

$$\begin{aligned} \text{st } \nu(B(x, r/2)) &= L(\nu)(B(x, r/2)) \\ &\leq L(\nu)(\varphi^{-1} B(\hat{x}, r)) \\ &= P(\nu)(B(\hat{x}, r)) \\ &= \mathcal{H}^\lambda(B(\hat{x}, r)) \end{aligned}$$

and similarly,

$$\begin{aligned} \text{st } \nu(B(x, r) \cap B(a, \rho)) &= L(\nu)(B(x, r) \cap B(a, \rho)) \\ &\geq L(\nu)(\varphi^{-1} B(\hat{x}, r) \cap \varphi^{-1} B(\hat{a}, \rho)) \\ &= P(\nu)(B(\hat{x}, r) \cap B(\hat{a}, \rho)) \\ &= \mathcal{H}^\lambda(B(\hat{x}, r) \cap B(\hat{a}, \rho)). \end{aligned}$$

Now we use the bounds given by Ahlfors regularity with the modified lower bound given by Lemma 9.2: If $x \in \mathfrak{X}$ and r, ρ are standard such that $r < R$, $r \leq \rho$ and $d(a, x) \leq \rho$,

$$k r^\lambda \leq \mathcal{H}^\lambda(B(\hat{x}, r) \cap B(\hat{a}, \rho))$$

We translate these inequalities as follows. Let $1/M$ be the density of ν . If $x \in \mathfrak{X}$ and r, ρ are standard such that $r < R$ and $d(a, x) \leq \rho$,

$$\text{st} \left(\frac{\text{card } B(x, r/2)}{M} \right) \leq K r^\lambda$$

and if in addition $r \leq \rho$,

$$\text{st} \left(\frac{\text{card}(B(x, r) \cap B(a, \rho))}{M} \right) \geq k r^\lambda.$$

Thus, for standard $K' > K$ and $k' < k$ and all standard ρ and standard r such that $0 < r < R$ and $x \in \mathfrak{X}$ such that $d(a, x) \leq \rho$

$$\text{card } B(x, r/2) \leq M K' r^\lambda$$

and if in addition $r \leq \rho$,

$$\text{card}(\mathbf{B}(x, r) \cap \mathbf{B}(a, \rho)) \geq M k' r^\lambda$$

Now the same inequalities holds for arbitrary $r \in {}^*\mathbb{R}$ for which $0 \ll r \ll R$ as is shown in the following lemma:

Lemma 9.4 *Suppose ψ is an internal nondecreasing function on ${}^*[0, \infty[$ and k, K are positive standard constants such that:*

$$k s^\lambda \leq \psi(s) \leq K s^\lambda \quad (21)$$

for all standard s in some interval $]\alpha, \beta[$ with α, β positive. Then inequality (21) holds for all $s \in {}^\mathbb{R}$ such that $\alpha \ll s \ll \beta$ with a possibly different value for K (not exceeding $(3/2)^\lambda$ the original value of K) and for k (but not less than $(2/3)^\lambda$ the original value of k).*

PROOF. If $\alpha \ll s \ll \beta$, there is a standard $r \ll \beta$ such that $s \leq r \leq 3/2 s$; for instance, take $r = 1/2 (\text{st}(s) + \min(\text{st}(\beta) + 3/2 \text{st}(s)))$ in case β is limited and $r = 5/4 \text{st}(s)$ otherwise. Thus

$$\psi(s) \leq \psi(r) \leq K r^\lambda \leq (3/2)^\lambda K s^\lambda,$$

and similarly for the lower bound. ■

Completion of Proof of Proposition of 9.3. By the lemma, for standard ρ , for $r \in {}^*\mathbb{R}$ such that $0 \ll r \ll R$ and $x \in \mathfrak{X}$ such that $d(a, x) \leq \rho$

$$\frac{\text{card} \mathbf{B}(x, r)}{r^\lambda} \leq M K' \quad (22)$$

and if $r \ll \rho$,

$$\frac{\text{card}(\mathbf{B}(x, r) \cap \mathbf{B}(a, \rho))}{r^\lambda} \geq M k' \quad (23)$$

For every $\rho \in \mathbb{N}$ let $r_\rho = 1/\rho$. Then

1. Inequality (22) holds for r such that $r_\rho \leq r \leq R - r_\rho$ and $d(a, x) \leq \rho$. In the case $R = +\infty$, inequality (22) holds for limited r such that $r_\rho \leq r$.
2. Inequality (23) holds for r which in addition satisfy $r \leq \rho - r_\rho$.

In particular, by saturation and overspill there is a $\rho \cong \infty$ and an $r_\infty \leq 1/\rho \cong 0$ such that these same inequalities hold for limited r such that $r_\infty \leq r \leq R - r_\infty$. Note that the additional restriction given by item (2) to insure the validity of inequality (23) disappears since r is limited.

Note that if $R = +\infty$, use overspill to conclude that these same inequalities hold for r such that $r_\infty \leq r \leq R_+$ for some unlimited R_+ .

To complete the proof, let $\mathcal{Y} = \mathbf{B}(a, \rho)$. ■

Remarks. Note that the only property that we have used of Hausdorff measure $\mathcal{H}^\lambda$ on (X, d) is that it satisfies an inequality of the form (10). Thus the same result is true of any Borel measure μ on X which is boundedly equivalent to λ -dimensional Hausdorff measure, that is which is absolutely continuous with respect to Hausdorff measure and for which the Radon-Nikodym derivative satisfies

$$c \leq \frac{d\mu}{d\mathcal{H}^\lambda}(x) \leq C \quad \text{for almost all } x \in X,$$

where $0 < c \leq C < \infty$ and “almost all” is relative to $\mathcal{H}^\lambda$.

10 Representation of Ahlfors Regular Spaces

Suppose δ is a hyperreal. An internal metric space is δ -connected iff for every $x, y \in \mathfrak{X}$ with $\delta \leq d(x, y)$, there is a sequence $x = x_0, \dots, x_n = y$ such that $d(x_i, x_{i+1}) \leq \delta$ for all $i \leq n-1$ and $n\delta \sim_O d(x, y)$. Gromov in [22] introduces a related notion called *long range connectedness*. Note that our definition is external.

Example 10.1 Suppose $(\mathfrak{X}, d)$ is an internal length space. Then for any $\delta > 0$, $(\mathfrak{X}, d)$ is δ -connected. Proof: Let $\delta \leq a = d(x, y)$. By assumption, there is an internal isometric map $f : [0, a] \rightarrow \mathfrak{X}$ such that $f(0) = x, f(a) = y$. Let n be the largest hyperinteger such that $(n-1)\delta < a$ and define $t_k = k\delta$ for $k \leq n-1$, $t_n = a$. As successive t_k 's differ by less than δ and f is an isometry, $d(f(t_{k+1}), f(t_k)) \leq \delta$. Moreover, $(n-1)\delta < a \leq n\delta$ so $1 \leq n\delta/a < 1 + \delta/a \leq 2$ and therefore $n\delta \sim_O a$.

We will use the following internal function between subsets of $\mathfrak{X}$: $d(A, B) = \sup_{x \in A} \inf_{y \in B} d(x, y)$. This function, unlike the Hausdorff distance function on subsets is not symmetric.

Proposition 10.1 Suppose $\mathcal{Y} \subseteq \mathfrak{X}$, and $d(\mathfrak{X}, \mathcal{Y}) \leq \delta$. If $\mathfrak{X}$ is δ -connected, then $\mathcal{Y}$ is 4δ -connected.

PROOF. Suppose $x, y \in \mathcal{Y}$ and $\delta < 4\delta \leq d(x, y)$. By assumption there is a sequence $x = x_0, \dots, x_n = y$ in $\mathfrak{X}$ such that $d(x_i, x_{i+1}) \leq \delta$ for all $0 \leq i \leq n-1$ and $n\delta \sim_O d(x, y)$. Since $d(\mathfrak{X}, \mathcal{Y}) \leq \delta$, there are $x = x'_0, x'_1, \dots, x'_{n-1}, x'_n = y$ in $\mathcal{Y}$ such that $d(x'_i, x_i) \leq 3/2\delta$ for $i = 1, \dots, n-1$. Therefore, for each i , $d(x'_i, x'_{i+1}) \leq d(x'_i, x_i) + d(x_i, x_{i+1}) + d(x_{i+1}, x'_{i+1}) \leq 4\delta$. ■

The δ -graph associated to a metric space $(\mathfrak{X}, d)$ is defined as follows; The vertices of the graph consists of the points of $\mathfrak{X}$ and edges $x \longleftrightarrow y$ iff $d(x, y) \leq \delta$. Do not confuse the δ -graph distance dist with $\delta \times \text{dist}$!

Proposition 10.2 Suppose $(\mathfrak{X}, d)$ is δ -connected and dist is the δ -graph distance on $\mathfrak{X}$. Then there is a $c \gg 0$ such that for all $x, y \in \mathfrak{X}$ with $\delta \leq d(x, y)$,

$$c\delta \text{dist}(x, y) \leq d(x, y) \leq \delta \text{dist}(x, y).$$

PROOF. If $\text{dist}(x, y) = n$, then by the definition of graph distance, there is a sequence $x = x_0, \dots, x_n = y$ such that $d(x_i, x_{i+1}) \leq \delta$. Thus by the triangle inequality, $d(x, y) \leq n\delta = \delta \text{dist}(x, y)$. In the other direction, suppose $\delta \leq d(x, y)$. $\text{dist}(x, y)$ is the smallest hyperinteger n such that there is a sequence $x = x_0, x_1, \dots, x_n = y$ in $\mathfrak{X}$ with $d(x_i, x_{i+1}) \leq \delta$ for all $0 \leq i < n - 1$. Since $(\mathfrak{X}, d)$ is a δ -space, there is a sequence with these properties and the additional property $0 \ll d(x, y)/(n\delta)$. In particular, $c_{x,y} = d(x, y)/(\text{dist}(x, y)\delta) \gg 0$. Take $c = \inf\{c_{x,y} : \delta \leq d(x, y)\} \gg 0$ which is the infimum over an internal condition. Thus $d(x, y) \geq c\delta \text{dist}(x, y)$. ■

Proposition 10.3 *Let $(\mathfrak{X}, d)$ be internal and δ -connected. Then there is a δ -separated $\mathcal{Y} \subseteq \mathfrak{X}$ with $d(\mathfrak{X}, \mathcal{Y}) \leq \delta$ satisfying the following property: If dist is the δ -graph metric on $\mathcal{Y}$, then δdist is S -Lipschitz equivalent to d on $\mathcal{Y}$.*

PROOF. Let $\mathcal{Y} \subseteq \mathfrak{X}$ be maximal δ -separated. In particular, $d(\mathfrak{X}, \mathcal{Y}) \leq \delta$. By Proposition 10.1, $\mathcal{Y}$ is 4δ -connected. Now if $x, y \in \mathcal{Y}$ with $x \neq y$, then $d(x, y) \geq \delta$ so by Proposition 10.2,

$$0 \ll \frac{d(x, y)}{4\delta \text{dist}(x, y)} \sim_O \frac{d(x, y)}{\delta \text{dist}(x, y)} \ll \infty.$$

Thus δdist on $\mathcal{Y}$ is S -Lipschitz equivalent to the metric d . ■

Note that nothing said so far implies $\mathcal{Y}$ can be taken to be hyperfinite.

Proposition 10.4 *Suppose $\delta \cong 0$ and $(\mathfrak{X}, d)$ is an internal δ -connected space such that all closed balls $\overline{B}(x, \delta)$ are $\ast$ -precompact. Then there is a graph $\mathcal{Y} \subseteq \mathfrak{X}$ with $d(\mathfrak{X}, \mathcal{Y}) \leq \delta$ such that δdist is S -Lipschitz equivalent to d and $\mathcal{Y}$ is hyperfinitely branching. In particular $B(a, r) \cap \mathcal{Y}$ is hyperfinite for any $r \in \ast \mathbb{R}$ and $a \in \mathcal{Y}$.*

PROOF. Let $\mathcal{Y} \subseteq \mathfrak{X}$ be as in Proposition 10.3 with the δ -graph structure. All nodes $a \in \mathcal{Y}$ have hyperfinitely many adjacent nodes. In fact let $A \subseteq \overline{B}(a, \delta)$ be hyperfinite and such that every $x \in \overline{B}(a, \delta)$ is distance $< \delta/2$ from some $x_A \in A$. Such a set A exists by $\ast$ -precompactness of $\overline{B}(a, \delta)$. By definition of the δ -graph, all nodes adjacent to a in the δ -graph of $\mathcal{Y}$ are members of $\overline{B}(a, \delta)$; since $\mathcal{Y}$ is δ -separated, all nodes in $\mathcal{Y}$ are at distance $\geq \delta$ from each other. Thus $x \mapsto x_A$ is injective on the set of nodes of $\mathcal{Y}$ adjacent to a . In particular the cardinality of this set is $\leq \text{card } A$ which is hyperfinite. Therefore $\mathcal{Y}$ is hyperfinitely branching. It follows that the set of points of $\mathcal{Y}$ at graph distance $< \infty$ (but possibly unlimited) from any $a \in \mathcal{Y}$ is hyperfinite. In particular, the set of points at distance $< \infty$ from any a is hyperfinite. ■

Remarks. The graph $\mathcal{Y}$ may have unlimited branching at each node. If $\mathfrak{X}$ has the property that $\overline{B}(x, \delta)$ is covered by a limited number of balls $B(x, \delta/2)$, then we may assume the graph $\mathcal{Y}$ has limited branching. For the set A in the above proof can be assumed limited.

Theorem 10.5 *Suppose (X, d) is an Ahlfors regular length space of dimension λ up to $R \in]0, \infty]$. Then there is a hyperfinite graph space $(\mathfrak{X}, d_{\mathfrak{X}})$, a Lipschitz equivalent metric d' on X and a hyperreal R_+ such that the following hold:*

1. (X, d') is a bounded component of $\diamond(\mathfrak{X}, d_{\mathfrak{X}})$.
2. For some uniform hypermeasure ν on $\mathfrak{X}$, $P_{\mathfrak{X}}(\nu)|_X = \mathcal{H}^\lambda$ (Hausdorff measure with respect to the original metric d , though see the remarks below).
3. $(\mathfrak{X}, d_{\mathfrak{X}})$ is of polynomial growth λ on $[R_-, R_+]$ where $R_- \cong 0$ and $0 \ll R_+ \ll \infty$ if R is finite or R_+ is an unlimited hyperreal in case $R = \infty$.

PROOF. Since (X, d) is a complete Ahlfors regular space, it is locally compact (See Proposition 6.1). By basic results on length spaces, all closed balls in (X, d) are compact. It follows $*(X, d)$ is an internal length space such that all closed balls are $*$ -compact. Moreover, by Proposition 2.8, the bounded component of X in $\diamond(*X)$ is X itself.

Let δ be an arbitrary positive infinitesimal. Consider the δ -graph structure on $*(X, d)$. Apply Proposition 10.4 to $\mathfrak{X} = *X$ to conclude the existence of a subgraph $\mathcal{Y}$ with the properties stated there. In particular, the metrics $\delta \text{ dist}$ and $*d$ on $\mathcal{Y}$ are S -Lipschitz equivalent. Since $d_{*X}(*X, \mathcal{Y}) \leq \delta$, it follows $\diamond \mathcal{Y} = \diamond(*X)$. Moreover, the metric $\diamond(\delta \text{ dist})$ is Lipschitz equivalent to $\diamond(*d)$. Now let $a \in \mathcal{Y}$ be such that $\hat{a} \in X$; then for r unlimited (but $r \in *\mathbb{R}$), $\mathcal{Z} = B(a, r) \cap \mathcal{Y}$ is hyperfinite, and such that $\diamond \mathcal{Z} \supseteq X$ and X is a bounded component of $\diamond \mathcal{Z}$.

Let d' be the metric $\diamond(\delta \text{ dist})$ restricted to X . By the previous paragraph, d' is Lipschitz equivalent to $\diamond(*d)|_X = d$. Now (X, d) is separable since it is a complete length space and in particular, X is σ -compact with respect to d' . Let μ be λ -dimensional Hausdorff measure relative to the metric d . μ is finite on compact sets and is boundedly equivalent to Hausdorff measure relative to the metric d' . It follows by Proposition 8.1 that there is a hypermeasure ν on $\mathcal{Z}$ such that $P_{\mathcal{Z}}(\nu)|_X = \mu$. Now $\text{supp}(\nu) = \{z \in \mathcal{Z} : \nu_z \neq 0\}$ is an internal set. In addition, by the Ahlfors regularity property of Hausdorff measure, $\diamond(\text{supp}(\nu)) \supseteq X$. Thus we can assume without loss of generality that $\nu_z \neq 0$ for all $z \in \mathcal{Z}$. By Proposition 8.4, we can also assume ν is a uniform hypermeasure on $\mathcal{Z}$ with density $1/M$. Applying Proposition 9.3 and the remarks after the proof of Proposition 9.3 to $(\mathcal{Z}, d_{\mathcal{Z}})$, we can immediately read off the main conclusion of the theorem. Thus there is a hyperfinite $\mathcal{Z}' \subseteq \mathcal{Z}$ such that $\diamond \mathcal{Z}' \supseteq X$ and $(\mathcal{Z}', d_{\mathcal{Z}})$ is of polynomial growth on some interval $[R_-, R]$ where $R_- \cong 0$. ■

Remarks. Note that in the previous result, $\mathcal{H}^\lambda$ may be replaced by any measure which is boundedly equivalent to $\mathcal{H}^\lambda$. See the remark after Proposition 9.3.

References

- [1] S. Albeverio, J.E. Fenstad, R. Höegh-Kron, and T. Lindström. *Nonstandard Analysis in Stochastic Analysis and Mathematical Physics*. Academic Press, New York, 1986.

- [2] H. Bass. The degree of polynomial growth of finitely generated nilpotent groups. *Proc. London Math. Soc.*, 25(3):603–614, 1972.
- [3] I. Chavel. *Riemannian Geometry: A Modern Introduction*. Cambridge Tracts in Mathematics. Cambridge University Press, 1993.
- [4] J. P. R. Christensen. On some measures analogous to Haar measure. *Math. Scand.*, 26:103–106, 1970.
- [5] G. David and S. Semmes. *Self-Similar Geometry through Metric and Measure*. Oxford Lecture Series in Mathematics. Clarendon Press, 1997.
- [6] J. Dieudonné. *Foundations of Modern Analysis*. Academic Press, 1960.
- [7] R. M. Dudley. *Real Analysis and Probability*. Wadsworth and Brooks/Cole, Pacific Grove, California, 1989.
- [8] K. J. Falconer. *The Geometry of Fractal Sets*. Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1985.
- [9] M. Gromov. Groups of polynomial growth and expanding maps. *Publications Mathématiques, I.H.E.S.*, 53, 1981.
- [10] M. Gromov. *Structures Métriques pour les Variétés Riemanniennes*. Textes mathématiques. Cedic/Fernand Nathan, 1981.
- [11] M. Gromov. *Metric Structures for Riemannian and Non-Riemannian Spaces*, volume 152 of *Progress in Mathematics*. Birkhäuser, 1998.
- [12] M. Hall. *Theory of Groups*. AMS Chelsea, Providence, second edition, 1976.
- [13] P. R. Halmos. *Measure Theory*. The University Series in Higher Mathematics. D. van Nostrand, Princeton, New Jersey, 1950.
- [14] C. W. Henson. Unbounded Loeb measures. *Proc. Amer. Math. Soc.*, 74:143–150, 1979.
- [15] A. Hurd and P.A. Loeb. *Introduction to Nonstandard Real Analysis*. Academic Press, New York, 1987.
- [16] Patrick Billingsley Jr. *Convergence of Probability Measures*. John Wiley and Sons, New York, 1968.
- [17] H.J. Keisler. *An Infinitesimal Approach to Stochastic Analysis*, volume 297 of *Memoirs, American Mathematical Society*. American Mathematical Society, 1984.
- [18] E. F. Krause. *Taxicab Geometry: An Adventure in Non-Euclidean Geometry*. Dover, New York, 1986.

- [19] P. Loeb. Conversion from nonstandard to standard measure spaces and applications in probability theory. *Trans. Amer. Math. Soc.*, 211:113–122, 1975.
- [20] P. Mattila. *Geometry of Sets and Measures in Euclidean Spaces*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1995.
- [21] F. Morgan. *Geometric Measure Theory: A Beginner's Guide*. Academic Press, Boston, 1988.
- [22] G. A. Niblo and M. A. Roller, editors. *Geometric Group Theory*, volume 182 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, U. K., 1993.
- [23] C. A. Rogers. *Hausdorff Measures*. Cambridge University Press, Cambridge, U. K., 1970.
- [24] L. van den Dries and A. J. Wilkie. Gromov's theorem on groups of polynomial growth and elementary logic. *J. of Algebra*, 89:349–374, 1984.

9112 Decatur Ave S.
 Bloomington, MN 55438
E-mail address: jt@mitre.org